



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/00, C07K 14/47, G01N 33/53	A2	(11) International Publication Number: WO 00/52151 (43) International Publication Date: 8 September 2000 (08.09.00)
(21) International Application Number: PCT/US00/05621 (22) International Filing Date: 3 March 2000 (03.03.00) (30) Priority Data: 60/123,117 5 March 1999 (05.03.99) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 60/123,117 (CIP) Filed on 5 March 1999 (05.03.99) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). AU-YOUNG, Janice [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US). LU, Dyung, Aina, M. [US/US]; 55 Park		Belmont Place, San Jose, CA 95136 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). (74) Agents: HAMLET-COX, Diana et al.; Incyte Pharmaceuticals, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: HUMAN SECRETORY PROTEINS (57) Abstract <p>The invention provides human secretory proteins (HSECP) and polynucleotides which identify and encode HSECP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of HSECP.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HUMAN SECRETORY PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of human secretory proteins
5 and to the use of these sequences in the diagnosis, treatment, and prevention of cancer, inflammation,
and gastrointestinal, cardiovascular, and neurological disorders.

BACKGROUND OF THE INVENTION

Protein transport and secretion are essential for cellular function. Protein transport is
10 mediated by a signal peptide located at the amino terminus of the protein to be transported or
secreted. The signal peptide is comprised of about ten to twenty hydrophobic amino acids which
target the nascent protein from the ribosome to a particular membrane bound compartment such as the
endoplasmic reticulum (ER). Proteins targeted to the ER may either proceed through the secretory
pathway or remain in any of the secretory organelles such as the ER, Golgi apparatus, or lysosomes.
15 Proteins that transit through the secretory pathway are either secreted into the extracellular space or
retained in the plasma membrane. Secreted proteins are often synthesized as inactive precursors that
are activated by post-translational processing events during transit through the secretory pathway.
Such events include glycosylation, proteolysis, and removal of the signal peptide by a signal
peptidase. Other events that may occur during protein transport include chaperone-dependent
20 unfolding and folding of the nascent protein and interaction of the protein with a receptor or pore
complex. Examples of secreted proteins with amino terminal signal peptides are discussed below and
include receptors, extracellular matrix molecules, cytokines, hormones, growth and differentiation
factors, neuropeptides, vasomediators, ion channels, transporters/pumps, and proteases. (Reviewed in
Alberts, B. et al. (1994) Molecular Biology of The Cell, Garland Publishing, New York, NY, pp. 557-
25 560, 582-592.)

G-protein coupled receptors (GPCRs) comprise a superfamily of integral membrane proteins
which transduce extracellular signals. Not all GPCRs contain N-terminal signal peptides. GPCRs
include receptors for biogenic amines such as dopamine, epinephrine, histamine, glutamate
(metabotropic-type), acetylcholine (muscarinic-type), and serotonin; for lipid mediators of
30 inflammation such as prostaglandins, platelet activating factor, and leukotrienes; for peptide
hormones such as calcitonin, C5a anaphylatoxin, follicle stimulating hormone, gonadotropin releasing
hormone, neurokinin, oxytocin, and thrombin; and for sensory signal mediators such as retinal
photopigments and olfactory stimulatory molecules. The structure of these highly conserved
receptors consists of seven hydrophobic transmembrane regions, cysteine disulfide bridges between
35 the second and third extracellular loops, an extracellular N-terminus, and a cytoplasmic C-terminus.

The N-terminus interacts with ligands, the disulfide bridges interact with agonists and antagonists, and the large third intracellular loop interacts with G proteins to activate second messengers such as cyclic AMP, phospholipase C, inositol triphosphate, or ion channels. (Reviewed in Watson, S. and Arkinstall, S. (1994) The G-protein Linked Receptor Facts Book, Academic Press, San Diego, CA, pp. 2-6; and Bolander, F.F. (1994) Molecular Endocrinology, Academic Press, San Diego, CA, pp. 162-176.)

Other types of receptors include cell surface antigens identified on leukocytic cells of the immune system. These antigens have been identified using systematic, monoclonal antibody (mAb)-based "shot gun" techniques. These techniques have resulted in the production of hundreds of mAbs directed against unknown cell surface leukocytic antigens. These antigens have been grouped into "clusters of differentiation" based on common immunocytochemical localization patterns in various differentiated and undifferentiated leukocytic cell types. Antigens in a given cluster are presumed to identify a single cell surface protein and are assigned a "cluster of differentiation" or "CD" designation. Some of the genes encoding proteins identified by CD antigens have been cloned and verified by standard molecular biology techniques. CD antigens have been characterized as both transmembrane proteins and cell surface proteins anchored to the plasma membrane via covalent attachment to fatty acid-containing glycolipids such as glycosylphosphatidylinositol (GPI). (Reviewed in Barclay, A. N. et al. (1995) The Leucocyte Antigen Facts Book, Academic Press, San Diego, CA, pp. 17-20.)

Matrix proteins (MPs) are transmembrane and extracellular proteins which function in formation, growth, remodeling, and maintenance of tissues and as important mediators and regulators of the inflammatory response. The expression and balance of MPs may be perturbed by biochemical changes that result from congenital, epigenetic, or infectious diseases. In addition, MPs affect leukocyte migration, proliferation, differentiation, and activation in the immune response. MPs are frequently characterized by the presence of one or more domains which may include collagen-like domains, EGF-like domains, immunoglobulin-like domains, and fibronectin-like domains. In addition, MPs may be heavily glycosylated and may contain an Arginine-Glycine-Aspartate (RGD) tripeptide motif which may play a role in adhesive interactions. MPs include extracellular proteins such as fibronectin, collagen, galectin, vitronectin and its proteolytic derivative somatomedin B; and cell adhesion receptors such as cell adhesion molecules (CAMs), cadherins, and integrins. (Reviewed in Ayad, S. et al. (1994) The Extracellular Matrix Facts Book, Academic Press, San Diego, CA, pp. 2-16; Ruoslahti, E. (1997) *Kidney Int.* 51:1413-1417; Sjaastad, M.D. and Nelson, W.J. (1997) *BioEssays* 19:47-55.)

Cytokines are secreted by hematopoietic cells in response to injury or infection. Interleukins, neurotrophins, growth factors, interferons, and chemokines all define cytokine families that work in

conjunction with cellular receptors to regulate cell proliferation and differentiation. In addition, cytokines effect activities such as leukocyte migration and function, hematopoietic cell proliferation, temperature regulation, acute response to infection, tissue remodeling, and apoptosis.

Chemokines, in particular, are small chemoattractant cytokines involved in inflammation, leukocyte proliferation and migration, angiogenesis and angiostasis, regulation of hematopoiesis, HIV infectivity, and stimulation of cytokine secretion. Chemokines generally contain 70-100 amino acids and are subdivided into four subfamilies based on the presence of conserved cysteine-based motifs. (Callard, R. and Gearing, A. (1994) The Cytokine Facts Book, Academic Press, New York, NY, pp. 181-190, 210-213, 223-227.)

Growth and differentiation factors are secreted proteins which function in intercellular communication. Some factors require oligomerization or association with MPs for activity. Complex interactions among these factors and their receptors trigger intracellular signal transduction pathways that stimulate or inhibit cell division, cell differentiation, cell signaling, and cell motility. Most growth and differentiation factors act on cells in their local environment (paracrine signaling). There are three broad classes of growth and differentiation factors. The first class includes the large polypeptide growth factors such as epidermal growth factor, fibroblast growth factor, transforming growth factor, insulin-like growth factor, and platelet-derived growth factor. The second class includes the hematopoietic growth factors such as the colony stimulating factors (CSFs). Hematopoietic growth factors stimulate the proliferation and differentiation of blood cells such as B-lymphocytes, T-lymphocytes, erythrocytes, platelets, eosinophils, basophils, neutrophils, macrophages, and their stem cell precursors. The third class includes small peptide factors such as bombesin, vasopressin, oxytocin, endothelin, transferrin, angiotensin II, vasoactive intestinal peptide, and bradykinin which function as hormones to regulate cellular functions other than proliferation.

Growth and differentiation factors play critical roles in neoplastic transformation of cells in vitro and in tumor progression in vivo. Inappropriate expression of growth factors by tumor cells may contribute to vascularization and metastasis of tumors. During hematopoiesis, growth factor misregulation can result in anemias, leukemias, and lymphomas. Certain growth factors such as interferon are cytotoxic to tumor cells both in vivo and in vitro. Moreover, some growth factors and growth factor receptors are related both structurally and functionally to oncoproteins. In addition, growth factors affect transcriptional regulation of both proto-oncogenes and oncosuppressor genes. (Reviewed in Pimentel, E. (1994) Handbook of Growth Factors, CRC Press, Ann Arbor, MI, pp. 1-9.)

Proteolytic enzymes or proteases either activate or deactivate proteins by hydrolyzing peptide bonds. Proteases are found in the cytosol, in membrane-bound compartments, and in the extracellular space. The major families are the zinc, serine, cysteine, thiol, and carboxyl proteases.

Ion channels, ion pumps, and transport proteins mediate the transport of molecules across

cellular membranes. Transport can occur by a passive, concentration-dependent mechanism or can be linked to an energy source such as ATP hydrolysis. Symporters and antiporters transport ions and small molecules such as amino acids, glucose, and drugs. Symporters transport molecules and ions unidirectionally, and antiporters transport molecules and ions bidirectionally. Transporter

5 superfamilies include facilitative transporters and active ATP-binding cassette transporters which are involved in multiple-drug resistance and the targeting of antigenic peptides to MHC Class I molecules. These transporters bind to a specific ion or other molecule and undergo a conformational change in order to transfer the ion or molecule across the membrane. (Reviewed in Alberts, B. et al. (1994) Molecular Biology of The Cell, Garland Publishing, New York, NY, pp. 523-546.)

10 Ion channels are formed by transmembrane proteins which create a lined passageway across the membrane through which water and ions, such as Na^+ , K^+ , Ca^{2+} , and Cl^- , enter and exit the cell. For example, chloride channels are involved in the regulation of the membrane electric potential as well as absorption and secretion of ions across the membrane. Chloride channels also regulate the internal pH of membrane-bound organelles.

15 Ion pumps are ATPases which actively maintain membrane gradients. Ion pumps are classified as P, V, or F according to their structure and function. All have one or more binding sites for ATP in their cytosolic domains. The P-class ion pumps include Ca^{2+} ATPase and Na^+/K^+ ATPase and function in transporting H^+ , Na^+ , K^+ , and Ca^{2+} ions. P-class pumps consist of two α and two β transmembrane subunits. The V- and F-class ion pumps have similar structures but transport only H^+ .
20 F class H^+ pumps mediate transport across the membranes of mitochondria and chloroplasts, while V-class H^+ pumps regulate acidity inside lysosomes, endosomes, and plant vacuoles.

A family of structurally related intrinsic membrane proteins known as facilitative glucose transporters catalyze the movement of glucose and other selected sugars across the plasma membrane. The proteins in this family contain a highly conserved, large transmembrane domain comprised of 12
25 α -helices, and several weakly conserved, cytoplasmic and exoplasmic domains. (Pessin, J. E., and Bell, G.I. (1992) *Annu. Rev. Physiol.* 54:911-930.)

Amino acid transport is mediated by Na^+ dependent amino acid transporters. These transporters are involved in gastrointestinal and renal uptake of dietary and cellular amino acids and in neuronal reuptake of neurotransmitters. Transport of cationic amino acids is mediated by the
30 system y+ family and the cationic amino acid transporter (CAT) family. Members of the CAT family share a high degree of sequence homology, and each contains 12-14 putative transmembrane domains. (Ito, K. and Groudine, M. (1997) *J. Biol. Chem.* 272:26780-26786.)

Hormones are secreted molecules that travel through the circulation and bind to specific receptors on the surface of, or within, target cells. Although they have diverse biochemical
35 compositions and mechanisms of action, hormones can be grouped into two categories. One category

includes small lipophilic hormones that diffuse through the plasma membrane of target cells, bind to cytosolic or nuclear receptors, and form a complex that alters gene expression. Examples of these molecules include retinoic acid, thyroxine, and the cholesterol-derived steroid hormones such as progesterone, estrogen, testosterone, cortisol, and aldosterone. The second category includes

- 5 hydrophilic hormones that function by binding to cell surface receptors that transduce signals across the plasma membrane. Examples of such hormones include amino acid derivatives such as catecholamines and peptide hormones such as glucagon, insulin, gastrin, secretin, cholecystokinin, adrenocorticotrophic hormone, follicle stimulating hormone, luteinizing hormone, thyroid stimulating hormone, and vasopressin. (See, for example, Lodish et al. (1995) Molecular Cell Biology, Scientific
10 American Books Inc., New York, NY, pp. 856-864.)

- Neuropeptides and vasomediators (NP/VM) comprise a large family of endogenous signaling molecules. Included in this family are neuropeptides and neuropeptide hormones such as bombesin, neuropeptide Y, neurotensin, neuromedin N, melanocortins, opioids, galanin, somatostatin, tachykinins, urotensin II and related peptides involved in smooth muscle stimulation, vasopressin,
15 vasoactive intestinal peptide, and circulatory system-borne signaling molecules such as angiotensin, complement, calcitonin, endothelins, formyl-methionyl peptides, glucagon, cholecystokinin and gastrin. NP/VMs can transduce signals directly, modulate the activity or release of other neurotransmitters and hormones, and act as catalytic enzymes in cascades. The effects of NP/VMs range from extremely brief to long-lasting. (Reviewed in Martin, C. R. et al. (1985) Endocrine
20 Physiology, Oxford University Press, New York, NY, pp. 57-62.)

The discovery of new human secretory proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cancer, inflammation, and gastrointestinal, cardiovascular, and neurological disorders.

25

SUMMARY OF THE INVENTION

- The invention features purified polypeptides, human secretory proteins, referred to collectively as "HSECP" and individually as "HSECP-1," "HSECP-2," "HSECP-3," "HSECP-4," "HSECP-5," "HSECP-6," "HSECP-7," "HSECP-8," "HSECP-9," "HSECP-10," "HSECP-11,"
30 "HSECP-12," "HSECP-13," "HSECP-14," "HSECP-15," "HSECP-16," "HSECP-17," "HSECP-18," "HSECP-19," "HSECP-20," "HSECP-21" and "HSECP-22." In one aspect, the invention provides an isolated polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, c) a biologically active
35 fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, or d) an

immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-22.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising
5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22. In one
10 alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:23-44.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group
15 consisting of SEQ ID NO:1-22, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, or d) an immunogenic fragment of an amino
20 acid sequence selected from the group consisting of SEQ ID NO:1-22. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a
30 polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID
35 NO:1-22.

The invention further provides an isolated polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:23-44, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:23-44, c) a polynucleotide sequence complementary to a), or d) a polynucleotide sequence complementary to b). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:23-44, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:23-44, c) a polynucleotide sequence complementary to a), or d) a polynucleotide sequence complementary to b). The method comprises a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 30 contiguous nucleotides. In another alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a pharmaceutical composition comprising an effective amount of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, and a pharmaceutically acceptable excipient. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional HSECP, comprising administering to a patient in need of such treatment the pharmaceutical composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22. The method comprises a) exposing a sample comprising the polypeptide to a

compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional HSECP,

5 comprising administering to a patient in need of such treatment the pharmaceutical composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, 10 c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an antagonist compound identified by 15 the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional HSECP, comprising administering to a patient in need of such treatment the pharmaceutical composition.

The invention further provides a method for screening a compound for effectiveness in 20 altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:23-44, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

25 BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding HSECP.

Table 2 shows features of each polypeptide sequence, including predicted signal peptides and 30 other motifs, and methods, algorithms, and searchable databases used for analysis of HSECP.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

35 Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones

encoding HSECP were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze HSECP, along with applicable descriptions, references, and threshold parameters.

5

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"HSECP" refers to the amino acid sequences of substantially purified HSECP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of HSECP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of HSECP either by directly interacting with HSECP or by acting on components of the biological pathway in which HSECP participates.

An "allelic variant" is an alternative form of the gene encoding HSECP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally-occurring form. Common mutational changes which give rise to

allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding HSECP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as HSECP or a polypeptide with at least one functional characteristic of HSECP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding HSECP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding HSECP. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent HSECP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of HSECP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where “amino acid sequence” is recited to refer to an amino acid sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

“Amplification” relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term “antagonist” refers to a molecule which inhibits or attenuates the biological activity of HSECP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of HSECP either by directly interacting with HSECP or by acting on components of the biological pathway in which HSECP participates.

The term “antibody” refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant.

Antibodies that bind HSECP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired.

- 5 Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to
10 immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense"
15 strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be
20 produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

25 The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic HSECP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of
30 polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization
35 between the nucleic acid strands. This is of particular importance in amplification reactions, which

depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding HSECP or fragments of HSECP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using the XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or more Incyte Clones and, in some cases, one or more public domain ESTs, using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that, when made, least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
	Ala	Gly, Ser
	Arg	His, Lys
25	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
30	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
	Lys	Arg, Gln, Glu
35	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
	Trp	Phe, Tyr
40	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

5 A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative
10 polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "fragment" is a unique portion of HSECP or the polynucleotide encoding HSECP which is
15 identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid
20 residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present
25 embodiments.

A fragment of SEQ ID NO:23-44 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:23-44, for example, as distinct from any other sequence in the same genome. A fragment of SEQ ID NO:23-44 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:23-44 from related
30 polynucleotide sequences. The precise length of a fragment of SEQ ID NO:23-44 and the region of SEQ ID NO:23-44 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-22 is encoded by a fragment of SEQ ID NO:23-44. A fragment of SEQ ID NO:1-22 comprises a region of unique amino acid sequence that specifically identifies
35 SEQ ID NO:1-22. For example, a fragment of SEQ ID NO:1-22 is useful as an immunogenic peptide

for the development of antibodies that specifically recognize SEQ ID NO:1-22. The precise length of a fragment of SEQ ID NO:1-22 and the region of SEQ ID NO:1-22 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

5 The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization
10 assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to
15 one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

20 The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

25 Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS
30 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequence pairs.

 Alternatively, a suite of commonly used and freely available sequence comparison algorithms
35 is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment

Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 11

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the hydrophobicity and acidity at the

site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term "humanized antibody" refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of identity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the

stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%.

Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune

disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

5 An "immunogenic fragment" is a polypeptide or oligopeptide fragment of HSECP which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of HSECP which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

10 The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate" refers to a change in the activity of HSECP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of HSECP.

15 The phrases "nucleic acid" or "nucleic acid sequence," as used herein, refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

20 "Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

25 "Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

30 "Probe" refers to nucleic acid sequences encoding HSECP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target
35 polynucleotide by complementary base-pairing. The primer may then be extended along the target

DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to

identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence.

5 This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter
10 sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

15 An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic
20 acids encoding HSECP, or fragments thereof, or HSECP itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or
25 synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

30 The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by
35 different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

5 "Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral
10 infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to
15 animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in
20 vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, and plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention
25 into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-
30 1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of
35 polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding

polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene
 5 between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having
 10 at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

15 THE INVENTION

The invention is based on the discovery of new human secretory proteins (HSECP), the polynucleotides encoding HSECP, and the use of these compositions for the diagnosis, treatment, or prevention of cancer, inflammation, and gastrointestinal, cardiovascular, and neurological disorders.

20 Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding HSECP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each HSECP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their
 25 corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. The Incyte clones in column 5 were used to assemble the consensus nucleotide sequence of each HSECP and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each
 30 polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; and column 6 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 6 were used to characterize each polypeptide through sequence homology and protein motifs. In column 5, the first line of each cell lists the amino acid
 35 residues comprising predicted signal peptide sequences located at the amino terminus of each

HSECP. Additional identifying motifs or signatures, such as a somatomedin B signature in SEQ ID NO:16 and seven putative transmembrane domains in SEQ ID NO:18, are also listed in column 5.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding HSECP. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:23-44 and to distinguish between SEQ ID NO:23-44 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express HSECP as a fraction of total tissues expressing HSECP. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing HSECP as a fraction of total tissues expressing HSECP. Column 5 lists the vectors used to subclone each cDNA library. In particular, three out of four cDNA libraries which express SEQ ID NO:23 are derived from cartilage and synovia associated with joint inflammation, and four out of five cDNA libraries which express SEQ ID NO:29 are derived from intestinal tissue. Furthermore, about half of the cDNA libraries expressing SEQ ID NO:34 are associated with inflammation or the hematopoietic/immune system. Likewise, about half of the cDNA libraries expressing SEQ ID NO:35 are associated with inflammation or the hematopoietic/immune system, and in particular, with inflammation of the joints. In addition, 82% of the cDNA libraries expressing SEQ ID NO:37 are derived from tissues of the nervous system. Finally, expression of SEQ ID NO:39 is detected solely in a subtracted prostate tumor cDNA library, and expression of SEQ ID NO:43 is detected only in two cDNA libraries derived from heart tissue.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding HSECP were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

The invention also encompasses HSECP variants. A preferred HSECP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the HSECP amino acid sequence, and which contains at least one functional or structural characteristic of HSECP.

The invention also encompasses polynucleotides which encode HSECP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:23-44, which encodes HSECP. The polynucleotide sequences of SEQ ID NO:23-44, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding HSECP. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding HSECP. A particular aspect of the invention encompasses a variant of a
5 polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:23-44 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:23-44. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of HSECP.

10 It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding HSECP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These
15 combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring HSECP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode HSECP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring HSECP under appropriately
20 selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding HSECP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the
25 nucleotide sequence encoding HSECP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode HSECP and HSECP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the
30 synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding HSECP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID
35 NO:23-44 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and

G33S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Perkin-Elmer). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Perkin-Elmer), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding HSECP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National

Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been
5 size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze
10 the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process
15 from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode HSECP may be cloned in recombinant DNA molecules that direct expression of
20 HSECP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express HSECP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter HSECP-encoding sequences for a variety of purposes including, but
25 not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

30 The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Crameri, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of HSECP, such as its biological or enzymatic activity or its ability
35 to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene

variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial"

5 breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable
10 manner.

In another embodiment, sequences encoding HSECP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser. 7*:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser. 7*:225-232.)

Alternatively, HSECP itself or a fragment thereof may be synthesized using chemical methods. For
15 example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of HSECP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

20 The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

25 In order to express a biologically active HSECP, the nucleotide sequences encoding HSECP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in
30 polynucleotide sequences encoding HSECP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding HSECP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding HSECP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional
35 transcriptional or translational control signals may be needed. However, in cases where only coding

sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used.

5 (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding HSECP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A

10 Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding HSECP. These include, but are not limited to, microorganisms such as bacteria transformed
15 with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

20 In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding HSECP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding HSECP can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORI plasmid (Life Technologies). Ligation of sequences encoding HSECP into the vector's multiple
25 cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509.) When large quantities of HSECP are needed, e.g. for the production of
30 antibodies, vectors which direct high level expression of HSECP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of HSECP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such
35 vectors direct either the secretion or intracellular retention of expressed proteins and enable

integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; and Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184.)

Plant systems may also be used for expression of HSECP. Transcription of sequences
5 encoding HSECP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105.)
10 These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding HSECP may be ligated into
15 an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses HSECP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-
20 based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.*
25 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of HSECP in cell lines is preferred. For example, sequences encoding HSECP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector.
30 Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

35 Any number of selection systems may be used to recover transformed cell lines. These

include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk⁻* and *apr⁻* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding HSECP is inserted within a marker gene sequence, transformed cells containing sequences encoding HSECP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding HSECP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding HSECP and that express HSECP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of HSECP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on HSECP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana

Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding HSECP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding HSECP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding HSECP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode HSECP may be designed to contain signal sequences which direct secretion of HSECP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding HSECP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric HSECP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of HSECP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST),

maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity
5 purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the HSECP encoding sequence and the heterologous protein sequence, so that HSECP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10).
10 A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled HSECP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the
15 T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

Fragments of HSECP may be produced not only by recombinant means, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be
20 achieved, for example, using the ABI 431A peptide synthesizer (Perkin-Elmer). Various fragments of HSECP may be synthesized separately and then combined to produce the full length molecule.

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of HSECP and human secretory proteins. In addition, the expression of HSECP is
25 closely associated with cancer, inflammation, and gastrointestinal, cardiovascular, and neurological disorders. Therefore, HSECP appears to play a role in cancer, inflammation, and gastrointestinal, cardiovascular, and neurological disorders. In the treatment of disorders associated with increased HSECP expression or activity, it is desirable to decrease the expression or activity of HSECP. In the treatment of disorders associated with decreased HSECP expression or activity, it is desirable to
30 increase the expression or activity of HSECP.

Therefore, in one embodiment, HSECP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HSECP. Examples of such disorders include, but are not limited to, a cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in
35 particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall

bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus ; an inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma,

5 atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's

10 thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal

15 circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a gastrointestinal disorder such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis,

20 pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic

25 encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha₁-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; a

30 cardiovascular disorder, and in particular, a disorder of the heart such as congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus

35 erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart

disease, congenital heart disease, and complications of cardiac transplantation; and a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural
5 muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-
10 Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic,
15 endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, and Tourette's disorder.

In another embodiment, a vector capable of expressing HSECP or a fragment or derivative
20 thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HSECP including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified HSECP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HSECP including, but not
25 limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of HSECP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HSECP including, but not limited to, those listed above.

In a further embodiment, an antagonist of HSECP may be administered to a subject to treat or
30 prevent a disorder associated with increased expression or activity of HSECP. Examples of such disorders include, but are not limited to, those cancer, inflammation, and gastrointestinal, cardiovascular, and neurological disorders described above. In one aspect, an antibody which specifically binds HSECP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express HSECP.

35 In an additional embodiment, a vector expressing the complement of the polynucleotide

encoding HSECP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of HSECP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of HSECP may be produced using methods which are generally known in the art. In particular, purified HSECP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind HSECP. Antibodies to HSECP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with HSECP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to HSECP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of HSECP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to HSECP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J.

Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and

Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce HSECP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g.,

10 Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

15 Antibody fragments which contain specific binding sites for HSECP may also be generated. For example, such fragments include, but are not limited to, $F(ab')_2$ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the $F(ab')_2$ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. 20 et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between HSECP and its 25 specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering HSECP epitopes is generally used, but a competitive binding assay may also be employed (Pound, *supra*).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for HSECP. Affinity is expressed as an 30 association constant, K_a , which is defined as the molar concentration of HSECP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple HSECP epitopes, represents the average affinity, or avidity, of the antibodies for HSECP. The K_a determined for a preparation of monoclonal antibodies, which are monospecific 35 for a particular HSECP epitope, represents a true measure of affinity. High-affinity antibody

preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the HSECP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of HSECP,

- 5 preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington, DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For
10 example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of HSECP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

- 15 In another embodiment of the invention, the polynucleotides encoding HSECP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding HSECP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding HSECP. Thus, complementary molecules or
20 fragments may be used to modulate HSECP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding HSECP.

- Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or
25 from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding HSECP. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

- Genes encoding HSECP can be turned off by transforming a cell or tissue with expression
30 vectors which express high levels of a polynucleotide, or fragment thereof, encoding HSECP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of
35 the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding HSECP. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may be employed.

- 5 Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco
10 NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

- Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example,
15 engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding HSECP.

- Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides,
20 corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

- Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared
25 by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding HSECP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA
30 constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

- RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase
35 linkages within the backbone of the molecule. This concept is inherent in the production of PNAs

and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

5 Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat.

10 Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

15 An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of HSECP, antibodies to HSECP, and mimetics, agonists, antagonists, or inhibitors of HSECP. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, 20 but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, 25 enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's 30 Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

35 Pharmaceutical preparations for oral use can be obtained through combining active

compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many

acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a
5 pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of HSECP, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein
10 the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of
15 administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example HSECP or fragments thereof, antibodies of HSECP, and agonists, antagonists or inhibitors of HSECP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined
20 by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and
25 animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the
30 subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4
35 days, every week, or biweekly depending on the half-life and clearance rate of the particular

formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art.

- 5 Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

- In another embodiment, antibodies which specifically bind HSECP may be used for the
- 10 diagnosis of disorders characterized by expression of HSECP, or in assays to monitor patients being treated with HSECP or agonists, antagonists, or inhibitors of HSECP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for HSECP include methods which utilize the antibody and a label to detect HSECP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or
- 15 without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring HSECP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of HSECP expression.

- 20 Normal or standard values for HSECP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to HSECP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of HSECP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values.
- 25 Deviation between standard and subject values establishes the parameters for diagnosing disease.

- In another embodiment of the invention, the polynucleotides encoding HSECP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of HSECP may be correlated
- 30 with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of HSECP, and to monitor regulation of HSECP levels during therapeutic intervention.

- In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding HSECP or closely related molecules may be used to identify nucleic acid sequences which encode HSECP. The specificity of the probe, whether it is
- 35 made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a

conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding HSECP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50%
5 sequence identity to any of the HSECP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:23-44 or from genomic sequences including promoters, enhancers, and introns of the HSECP gene.

Means for producing specific hybridization probes for DNAs encoding HSECP include the cloning of polynucleotide sequences encoding HSECP or HSECP derivatives into vectors for the
10 production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ³²P or ³⁵S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

15 Polynucleotide sequences encoding HSECP may be used for the diagnosis of disorders associated with expression of HSECP. Examples of such disorders include, but are not limited to, a cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary,
20 pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact
25 dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis,
30 psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a gastrointestinal disorder such as dysphagia, peptic esophagitis, esophageal spasm, esophageal
35 stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea,

emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis,

5 Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha₁-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein

10 obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; a cardiovascular disorder, and in particular, a disorder of the heart such as congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease,

15 degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation; and a neurological

20 disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess,

25 suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central

30 nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia,

35 catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic

neuralgia, and Tourette's disorder. The polynucleotide sequences encoding HSECP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered HSECP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding HSECP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding HSECP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding HSECP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of HSECP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding HSECP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding HSECP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding HSECP, or a fragment of a polynucleotide complementary to the polynucleotide encoding HSECP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

Methods which may also be used to quantify the expression of HSECP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding HSECP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the

Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding HSECP on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, HSECP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between HSECP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with HSECP, or fragments thereof, and washed. Bound HSECP is then detected by methods well known in the art. Purified HSECP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding HSECP specifically compete with a test compound for binding HSECP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with HSECP.

In additional embodiments, the nucleotide sequences which encode HSECP may be used in

any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding
5 description, utilize the present invention to its fullest extent. The following preferred specific
embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder
of the disclosure in any way whatsoever.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific
10 embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/123,117, are hereby expressly incorporated by reference.

15 **EXAMPLES**

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a
20 monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-

1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSORT1 plasmid (Life Technologies), pcDNA2.1 plasmid (Invitrogen, Carlsbad CA), or pINCY plasmid (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

cDNA sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Perkin-Elmer) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Perkin-Elmer) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:23-44. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs

from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding HSECP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Extension of HSECP Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:23-44 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR

was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM

BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

In like manner, the nucleotide sequences of SEQ ID NO:23-44 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

5 VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:23-44 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06
10 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ - 32 P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based
15 hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature
20 under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array
25 elements on the surface of a substrate. (See, e.g., Baldeschweiler, *supra*.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and
30 patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software
35 well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or

fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

Sequences complementary to the HSECP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring HSECP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of HSECP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the HSECP-encoding transcript.

IX. Expression of HSECP

Expression and purification of HSECP is achieved using bacterial or virus-based expression systems. For expression of HSECP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3).

Antibiotic resistant bacteria express HSECP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of HSECP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding HSECP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945.)

In most expression systems, HSECP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from HSECP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified HSECP obtained by these methods can be used directly in the following activity assay.

X. Demonstration of HSECP Activity

An assay for HSECP activity measures the expression of HSECP on the cell surface. cDNA encoding HSECP is subcloned into an appropriate mammalian expression vector suitable for high levels of cDNA expression. The resulting construct is transfected into a nonhuman cell line such as NIH3T3. Cell surface proteins are labeled with biotin using methods known in the art. Immunoprecipitations are performed using HSECP-specific antibodies, and immunoprecipitated samples are analyzed using SDS-PAGE and immunoblotting techniques. The ratio of labeled immunoprecipitant to unlabeled immunoprecipitant is proportional to the amount of HSECP expressed on the cell surface.

Alternatively, an assay for HSECP activity measures the amount of HSECP in secretory, membrane-bound organelles. Transfected cells as described above are harvested and lysed. The lysate is fractionated using methods known to those of skill in the art, for example, sucrose gradient ultracentrifugation. Such methods allow the isolation of subcellular components such as the Golgi apparatus, ER, small membrane-bound vesicles, and other secretory organelles.

Immunoprecipitations from fractionated and total cell lysates are performed using HSECP-specific antibodies, and immunoprecipitated samples are analyzed using SDS-PAGE and immunoblotting techniques. The concentration of HSECP in secretory organelles relative to HSECP in total cell lysate is proportional to the amount of HSECP in transit through the secretory pathway.

XI. Functional Assays

HSECP function is assessed by expressing the sequences encoding HSECP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCR3.1

plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of HSECP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding HSECP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding HSECP and other genes of interest can be analyzed by northern analysis or microarray techniques.

XII. Production of HSECP Specific Antibodies

HSECP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the HSECP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A

peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-HSECP activity by, for example, binding the peptide or HSECP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIII. Purification of Naturally Occurring HSECP Using Specific Antibodies

Naturally occurring or recombinant HSECP is substantially purified by immunoaffinity chromatography using antibodies specific for HSECP. An immunoaffinity column is constructed by covalently coupling anti-HSECP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing HSECP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of HSECP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/HSECP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and HSECP is collected.

XIV. Identification of Molecules Which Interact with HSECP

HSECP, or biologically active fragments thereof, are labeled with ^{125}I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled HSECP, washed, and any wells with labeled HSECP complex are assayed. Data obtained using different concentrations of HSECP are used to calculate values for the number, affinity, and association of HSECP with the candidate molecules.

Alternatively, molecules interacting with HSECP are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, *Nature* 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	23	078811	SYNORAB01	078811H1 (SYNORAB01), 078811R6 (SYNORAB01), 078811T6 (SYNORAB01),
2	24	371156	LUNGNOT02	077182F1 (SYNORAB01), 077182R1 (SYNORAB01), 289599H1 (TMLR3DT01), 371156H1 (LUNGNOT02), 523415H1 (MMLR2DT01), 3438708F6 (PENCNOT06), 3772179F6 (BRSTNOT25), 3808004H1 (CONTTUT01)
3	25	584050	PROSNOT02	584050H1 (PROSNOT02), 1726563T6 (PROSNOT14), 1739666R6 (HIPONON01), 1856214F6 (PROSNOT18), 2305379R6 (NGANNOT01), 2681374F6 (SINIUCT01), 4070575H1 (KIDNNOT26), 5274539H1 (OVARINOT02), SZA101719F1, SASA02714F1
4	26	863808	BRAITUT03	851821H1 (NGANNOT01), 863808H1 (BRAITUT03), 863808T1 (BRAITUT03), 2735728F6 (OVARNOT09)
5	27	978433	BRSTNOT02	054823R1 (FIBRNOT01), 978433H1 (BRSTNOT02), 978433R1 (BRSTNOT02), 1867687T6 (SKINBIT01), 2503122H1 (CONUTUT01), 2532586H1 (GBLANOT02), 3411659H1 (BRSTTUS08)
6	28	1655369	PROSTUT08	746013R1 (BRAITUT01), 944864R6 (RATRNOT02), 1539790R1 (SINTTUT01), 1617847F6 (BRAITUT12), 1655369F6 (PROSTUT08), 1655369H1 (PROSTUT08), 1673290F6 (BLADNOT05), 2056840R6 (BEPINOT01), 3407992H1 (PROSTUS08), 4077365H1 (PANCNOT19), 4098012H1 (BRAITUT26)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
7	29	1703244	DUODNOT02	273878H1 (PANCIT03), 1632051H1 (COLNNOT19), 1703244H1 (DUODNOT02), 4176224H1 (SINTNOT21)
8	30	1730819	BRSTTUT08	862188R1 (BRAITUT03), 1399644F1 (BRAITUT08), 1443690F1 (THYRNOT03), 1596446F1 (BRAINOT14), 1730819F6 (BRSTTUT08), 1730819H1 (BRSTTUT08), 2304942H1 (NGANNOT01), 2868843H1 (THYRNOT10), 3395457H1 (LUNGNOT28)
9	31	1757161	PITUNOT03	864626R1 (BRAITUT03), 1231577H1 (BRAITUT01), 1395593F1 (THYRNOT03), 1466766T1 (PANCUTUT02), 1466766T6 (PANCUTUT02), 1597583F6 (BRAINOT14), 1757161H1 (PITUNOT03), 1757161R6 (PITUNOT03), 1757161T6 (PITUNOT03)
10	32	1976095	PANCUTUT02	864976T1 (BRAITUT03), 1976095H1 (PANCUTUT02), 4616101H1 (BRAYDIT01), SARB01143F1, SARB01861F1, SAJA02355F1
11	33	2169991	ENDCNOT03	2169991H1 (ENDCNOT03), 2313548R3 (NGANNOT01), 2727735T3 (OVARUT05), 3095616H1 (CERVNOT03), 3186421H1 (THYMNON04), 3493007H1 (ADRETUT07), SASA03601F1, SBDA00157F1
12	34	2616827	GBLANOT01	2616827H1 (GBLANOT01), 4637616F6 (MYEPTXT01), 5219252H1 (BRSTNOT35)
13	35	2991370	KIDNFET02	080447R1 (SYNORAB01), 190292R1 (SYNORAB01), 192279R1 (SYNORAB01), 2724873H1 (OVARUTUT05), 2991370H1 (KIDNFET02), SAUA03596F1, SAUA01525F1

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
14	36	3031062	TYMNOT05	1354027T3 (LUNGNOT09), 2889215F6 (LUNGFET04), 3031062H1 (TYMNOT05), SBAA04235F1, SBAA0620F1, SBAA01760F1, SBAA04365F2, SBAA01589F1
15	37	3101617	BRAINOT20	3101617H1 (BRAINOT20), 3335717F6 (BRAIFET01), SCAA04574V1, SCAA04351V1, SCAA03628V1, SCAA05459V1, SCAA01004V1
16	38	3216178	TESTNOT07	632084R6 (KIDNNOT05), 1993593T6 (CORPNOT02), 3216178F6 (TESTNOT07), 3216178H1 (TESTNOT07), 4914242H1 (LIVRFET05)
17	39	3406803	PROSTUS08	3406803F6 (PROSTUS08), 3406803H1 (PROSTUS08), 3406803T6 (PROSTUS08)
18	40	3468066	BRAIDIT01	659544H1 (BRAINOT03), 897142R1 (BRSTNOT05), 1321038F1 (BLADNOT04), 1351888F1 (LATRTUT02), 1485695F1 (CORPNOT02), 1507666F1 (LUNGNOT14), 2953291H1 (KIDNFET01), 3468066H1 (BRAIDIT01), 4426018F6 (BRAPDIT01)
19	41	3592862	293TF5T01	941610H1 (ADRENOT03), 1288036H1 (BRAINOT11), 1687969F6 (PROSTUT10), 3592862H1 (293TF5T01)
20	42	3669422	KIDNTUT16	3669422F6 (KIDNTUT16), 3669422H1 (KIDNTUT16), 3669422T6 (KIDNTUT16), 5445503H1 (LNODNOT12)
21	43	3688740	HEAANOT01	462098R6 (LATRNOT01), 3688740H1 (HEAANOT01)
22	44	3742589	THYMNOT08	938795R1 (CERVNOT01), 1255960T1 (MENITUT03), 1530330R1 (PANCNOT04), 3742589H1 (THYMNOT08), SBQA04142D1

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Potential Signal Peptides and Other Signature Sequences	Analytical Methods
1	182	S149 T151 S13		M1-S16; M1-P18	MOTIFS SPSCAN HMM
2	125	T87 S36 T78 T111	N75	M1-S37	MOTIFS SPSCAN HMM
3	320	S21 T63 S267 S300 T164	N40	M1-G20; M1-G23	MOTIFS SPSCAN HMM
4	234	T74 S198 T210 T227 S131 T195		M1-A30; M1-G25	MOTIFS SPSCAN HMM
5	278	S64 S132 S230 T252 S7 S179	N221	M1-A65	MOTIFS SPSCAN HMM
6	136	S98 S99		M1-P54	MOTIFS SPSCAN HMM
7	109	T57		M1-T20	MOTIFS SPSCAN HMM

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Potential Signal Peptides and Other Signature Sequences	Analytical Methods
8	262	S73 S91 S136 S86	N182	M1-A25; M1-G27	MOTIFS SPSCAN HMM
9	384	S11 T140 S32 S185 S232 S306 S378	N50 N59 N62 N304	M1-G31 Transmembrane domains: G195-A220 L74-F91 A142-T160	MOTIFS SPSCAN HMM
10	244	T95 S110 T208 T44 S47 S53 S69 S152 T194		M1-A23	MOTIFS SPSCAN HMM
11	326	S138 T59 S239	N129 N237	M1-G40	MOTIFS SPSCAN HMM
12	105	S96 T24	N94	M1-A19; M1-A23	MOTIFS SPSCAN HMM

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Potential Signal Peptides and Other Signature Sequences	Analytical Methods
13	626	S188 S593 T61 T118 T144 S252 T275 T410 S423 S501 T506 S524 T536 S550 S28 T146 S336 T467 S583 T588		M1-G27 RGD tripeptide: R271-D273	MOTIFS SPSCAN HMM
14	296	S205 T295 S109 T165 T214 S244 S73 S225 Y236	N40 N53 N204 N281	M1-G23	MOTIFS SPSCAN HMM
15	249	S7 T123 S233 S237 T151	N100	M1-G23; M1-A22 Transmembrane domains: T175-A193 P97-F120	MOTIFS SPSCAN HMM
16	124	S62 S109		M1-G37 Somatomedin B signature: L36-S105	MOTIFS SPSCAN HMM PROFILESCAN
17	101	T59 S67		M1-S40; M1-S28	MOTIFS SPSCAN HMM

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Potential Signal Peptides and Other Signature Sequences	Analytical Methods
18	540	S42 S38 T80 T172 S179 S326 S519 S531 T131 T250 T278 Y99	N78 N88 N170 N347 N448 N457	M1-C29; M1-I31 Transmembrane domains: V307-N327 L411-I428 A140-L163 D366-I385 Y99-Y122 F496-C513 I56-Y73	MOTIFS SPSCAN HMM PRINTS
19	108	S34 S64 S29 S47 Y96		M1-R22; M1-S23	MOTIFS SPSCAN HMM
20	114	S108 S5 S73 T85		M1-T32	MOTIFS SPSCAN HMM
21	114	T8	N6	M1-A34	MOTIFS SPSCAN HMM
22	287	S25 T75 T134 T139 S225 T255 S254		M1-P20; M1-A22 RGD tripeptide: R172-D174	MOTIFS SPSCAN HMM

Table 3

Nucleotide SEQ ID NO:	Selected Fragment(s)	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
23	110-154 542-586	Musculoskeletal (0.750) Reproductive (0.250)	Inflammation (0.750) Cancer (0.250)	PBLUESCRIPT
24	230-271	Reproductive (0.333) Musculoskeletal (0.190) Cardiovascular (0.143)	Cancer (0.476) Inflammation (0.333)	PBLUESCRIPT
25	109-153 649-693	Reproductive (0.314) Nervous (0.235) Gastrointestinal (0.157)	Cancer (0.608) Inflammation (0.196) Cell Proliferation (0.118)	PSPORT1
26	116-160	Reproductive (0.333) Nervous (0.178) Cardiovascular (0.156)	Cancer (0.600) Cell Proliferation (0.178) Inflammation (0.133)	PSPORT1
27	228-272	Reproductive (0.333) Cardiovascular (0.244) Gastrointestinal (0.111)	Cancer (0.667) Cell Proliferation (0.133) Inflammation (0.089)	PSPORT1
28	1945-1989	Nervous (0.301) Reproductive (0.219) Gastrointestinal (0.137)	Cancer (0.452) Inflammation (0.205) Trauma (0.164)	pINCY
29	271-315	Gastrointestinal (0.800) Nervous (0.200)	Trauma (0.600) Cancer (0.200) Inflammation (0.200)	pINCY
30	218-262	Reproductive (0.249) Nervous (0.195) Gastrointestinal (0.136)	Cancer (0.521) Inflammation (0.207) Cell Proliferation (0.172)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragment(s)	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
31	1190-1234	Nervous (0.324) Reproductive (0.250) Hematopoietic/Immune (0.118)	Cancer (0.426) Inflammation (0.279) Cell Proliferation (0.147)	PSPORT1
32	487-531	Gastrointestinal (0.333) Nervous (0.259) Reproductive (0.111)	Cancer (0.593) Inflammation (0.148) Cell Proliferation (0.111)	pINCY
33	1513-1557	Reproductive (0.255) Nervous (0.216) Hematopoietic/Immune (0.176)	Cancer (0.373) Inflammation (0.255) Cell Proliferation (0.196)	pINCY
34	270-317	Hematopoietic/Immune (0.455) Musculoskeletal (0.182) Nervous (0.182)	Inflammation (0.545) Cancer (0.273) Cell Proliferation (0.091)	pINCY
35	1299-1343 1956-2000	Musculoskeletal (0.519) Nervous (0.148) Cardiovascular (0.111)	Inflammation (0.481) Cancer (0.296) Trauma (0.074)	pINCY
36	651-695	Nervous (0.250) Developmental (0.167) Gastrointestinal (0.167)	Cell Proliferation (0.667) Inflammation (0.250) Cancer (0.167)	pINCY
37	218-262	Nervous (0.818) Gastrointestinal (0.091) Reproductive (0.091)	Cancer (0.545) Cell Proliferation (0.091) Inflammation (0.091)	pINCY
38	290-334	Hematopoietic/Immune (0.250) Urologic (0.250) Developmental (0.125)	Cell Proliferation (0.375) Inflammation (0.250) Cancer (0.125)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragment(s)	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
39	434-478	Reproductive (1.000)	Cancer (1.000)	pINCY
40	326-370	Nervous (0.242) Reproductive (0.220) Gastrointestinal (0.121)	Cancer (0.462) Inflammation (0.280) Cell Proliferation (0.121)	pINCY
41	165-209	Nervous (0.333) Gastrointestinal (0.200) Cardiovascular (0.133)	Cancer (0.467) Cell Proliferation (0.200) Inflammation (0.133)	pINCY
42	273-317	Hematopoietic/Immune (0.312) Nervous (0.188) Reproductive (0.167)	Cancer (0.354) Inflammation (0.312) Cell Proliferation (0.146)	pINCY
43	273-317	Cardiovascular (1.000)	Inflammation (0.500)	pINCY
44	435-479	Reproductive (0.297) Nervous (0.217) Gastrointestinal (0.109)	Cancer (0.464) Cell Proliferation (0.196) Inflammation (0.167)	pINCY

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
23	SYNORAB01	This library was constructed using RNA isolated from the synovial membrane tissue of a 68-year-old Caucasian female with rheumatoid arthritis.
24	LUNGNOT02	This library was constructed using RNA isolated from the lung tissue of a 47-year-old Caucasian male who died of a subarachnoid hemorrhage.
25	PROSNOT02	This library was constructed using RNA isolated from diseased prostate tissue removed from a 50-year-old Caucasian male during a retropubic prostatectomy. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated adenocarcinoma Gleason grade 3+3. Patient history included dysuria, carcinoma in situ of prostate, coronary atherosclerosis, and hyperlipidemia.
26	BRAITUT03	This library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe of a 17-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a grade 4 fibrillary giant and small-cell astrocytoma. Family history included benign hypertension and cerebrovascular disease.
27	BRSTNOT02	This library was constructed using RNA isolated from diseased breast tissue removed from a 55-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated proliferative fibrocystic changes characterized by apocrine metaplasia, sclerosing adenosis, cyst formation, and ductal hyperplasia without atypia. Pathology for the associated tumor tissue indicated an invasive grade 4 mammary adenocarcinoma. Patient history included atrial tachycardia and a benign neoplasm. Family history included cardiovascular and cerebrovascular disease.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
28	PROSTUT08	This library was constructed using RNA isolated from prostate tumor tissue removed from a 60-year-old Caucasian male during radical prostatectomy and regional lymph node excision. Pathology indicated an adenocarcinoma (Gleason grade 3+4). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA). Patient history included a kidney cyst, and hematuria. Family history included tuberculosis, cerebrovascular disease, and arteriosclerotic coronary artery disease.
29	DUODNOT02	This library was constructed using RNA isolated from duodenal tissue of an 8-year-old Caucasian female, who died from head trauma. Serology was positive for cytomegalovirus (CMV).
30	BRSTTUT08	This library was constructed using RNA isolated from breast tumor tissue removed from a 45-year-old Caucasian female during unilateral extended simple mastectomy. Pathology indicated invasive nuclear grade 2-3 adenocarcinoma, ductal type, with 3 of 23 lymph nodes positive for metastatic disease. Greater than 50% of the tumor volume was in situ, both comedo and non-comedo types. Immunostains were positive for estrogen/progesterone receptors, and uninvolved tissue showed proliferative changes. The patient concurrently underwent a total abdominal hysterectomy. Patient history included valvuloplasty of mitral valve without replacement, rheumatic mitral insufficiency, and rheumatic heart disease. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.
31	PITUNOT03	This library was constructed using RNA isolated from pituitary tissue of a 46-year-old Caucasian male who died from colon cancer.
32	PANCTUT02	This library was constructed using RNA isolated from pancreatic tumor tissue removed from a 45-year-old Caucasian female during radical pancreaticoduodenectomy. Pathology indicated a grade 4 anaplastic carcinoma. Family history included benign hypertension, hyperlipidemia and atherosclerotic coronary artery disease.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
33	ENDCNOT03	This library was constructed using RNA isolated from dermal microvascular endothelial cells removed from a neonatal Caucasian male.
34	GBLANOT01	This library was constructed using RNA isolated from diseased gallbladder tissue removed from a 53-year-old Caucasian female during a cholecystectomy. Pathology indicated mild chronic cholecystitis and cholelithiasis with approximately 150 mixed gallstones. Family history included benign hypertension.
35	KIDNFET02	This library was constructed using RNA isolated from kidney tissue removed from a Caucasian male fetus who was stillborn with a hypoplastic left heart at 23 weeks' gestation.
36	TLYMNOT05	This library was constructed using RNA isolated from nonactivated Th2 cells. The cells were differentiated from umbilical cord CD4 T cells with IL-4 in the presence of anti-IL-12 antibodies and B7-transfected COS cells.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
37	BRAINOT20	<p>This library was constructed using RNA isolated from diseased brain tissue removed from the left temporal lobe of a 27-year-old Caucasian male during a brain lobectomy. Pathology for the left temporal lobe, including the mesial temporal structures, indicated focal, marked pyramidal cell loss and gliosis in hippocampal sector CA1, consistent with mesial temporal sclerosis. The left frontal lobe showed a focal deep white matter lesion, characterized by marked gliosis, calcifications, and hemosiderin-laden macrophages, consistent with a remote perinatal injury. This frontal lobe tissue also showed mild to moderate generalized gliosis, predominantly subpial and subcortical, consistent with chronic seizure disorder. GFAP was positive for astrocytes. Family history included brain cancer.</p>
38	TESTNOT07	<p>This library was constructed using RNA isolated from testicular tissue removed from a 31-year-old Caucasian male during a unilateral orchiectomy (excision of testis). Pathology indicated a mass containing a large subcapsular hematoma with laceration of the tunica albuginea. The surrounding testicular parenchyma was extensively necrotic. The patient presented with a trunk injury.</p>
39	PROSTUS08	<p>This subtracted library was constructed using 2.36 million clones from a prostate tumor library and was subjected to one round of subtractive hybridization with 448,000 clones from a prostate library. The starting library for subtraction was constructed using RNA isolated from a prostate tumor removed from a 59-year-old Caucasian male during a radical prostatectomy with regional lymph node excision. Pathology indicated adenocarcinoma (Gleason grade 3+3) and adenofibromatous hyperplasia. The patient presented with elevated prostate-specific antigen (PSA). Patient history included colon diverticuli, asbestosis, and thrombophlebitis. Family history included multiple myeloma, hyperlipidemia, and rheumatoid arthritis. Subtractive hybridization conditions were based on the methodologies of Swaroop et al., NAR (1991) 19:1954 and Bonaldo, et al. Genome Research (1996) 6:791.</p>
40	BRAIDT01	<p>This library was constructed using RNA isolated from diseased brain tissue. Patient history included multiple sclerosis, type II lesion.</p>

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
41	293TF5T01	This library was constructed using RNA isolated from a transformed embryonal cell line (293-EBNA) derived from kidney epithelial tissue transfected with bgal. The cells were transformed with adenovirus 5 DNA.
42	KIDNTUT16	This library was constructed using RNA isolated from left pole kidney tumor tissue removed from a 53-year-old Caucasian female during a nephroureterectomy. Pathology indicated grade 2 renal cell carcinoma. Patient history included hyperlipidemia, cardiac dysrhythmia, menorrhagia, cerebrovascular disease, atherosclerotic coronary artery disease, and tobacco abuse. Family history included cerebrovascular disease and atherosclerotic coronary artery disease.
43	HEANOT01	This library was constructed using RNA isolated from right coronary and right circumflex coronary artery tissue removed from the explanted heart of a 46-year-old Caucasian male during a heart transplantation. Patient history included myocardial infarction from total occlusion of the left anterior descending coronary artery, atherosclerotic coronary artery disease, hyperlipidemia, myocardial ischemia, dilated cardiomyopathy, left ventricular dysfunction, and tobacco abuse. Previous surgeries included cardiac catheterization. Family history included atherosclerotic coronary artery disease.
44	THYMNOT08	This library was constructed using RNA isolated from thymus tissue removed from a 4-month-old Caucasian male during a total thymectomy and open heart repair of atrioventricular canal defect using hypothermia. Patient presented with a congenital heart anomaly, congestive heart failure, and Down syndrome. Patient history included abnormal thyroid function study, premature birth, and right and left heart angiocardiology.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfilesScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality score \geq GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <i>supra</i> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- 5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-22,
 b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-22,
 c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22, and
10 d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-22.

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-22.

15

3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide of claim 3 selected from the group consisting of SEQ ID NO:23-44.

20

5. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

6. A cell transformed with a recombinant polynucleotide of claim 5.

25

7. A transgenic organism comprising a recombinant polynucleotide of claim 5.

8. A method for producing a polypeptide of claim 1, the method comprising:

- 30 a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
 b) recovering the polypeptide so expressed.

35

9. An isolated antibody which specifically binds to a polypeptide of claim 1.

10. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:23-44,
- b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:23-44,
- c) a polynucleotide sequence complementary to a),
- d) a polynucleotide sequence complementary to b), and
- e) an RNA equivalent of a)-d).

11. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 10.

12. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 10, the method comprising:

- a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

13. A method of claim 12, wherein the probe comprises at least 30 contiguous nucleotides.

14. A method of claim 12, wherein the probe comprises at least 60 contiguous nucleotides.

15. A pharmaceutical composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

16. A method for treating a disease or condition associated with decreased expression of functional HSECP, comprising administering to a patient in need of such treatment the pharmaceutical composition of claim 15.

17. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and

b) detecting agonist activity in the sample.

18. A pharmaceutical composition comprising an agonist compound identified by a method of claim 17 and a pharmaceutically acceptable excipient.

5

19. A method for treating a disease or condition associated with decreased expression of functional HSECP, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 18.

10

20. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

15

21. A pharmaceutical composition comprising an antagonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

20

22. A method for treating a disease or condition associated with overexpression of functional HSECP, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 21.

23. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 4, the method comprising:

25

- a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.

TANG, Y. Tom

LAL, Preeti

BAUGHN, Mariah R.

YUE, Henry

AU-YOUNG, Janice

LU, Dyung Aina M.

AZIMZAI, Yalda

<120> HUMAN SECRETORY PROTEINS

<130> PF-0675 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/123,117

<151> 1999-03-05

<160> 44

<170> PERL Program

<210> 1

<211> 182

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 078811CD1

<400> 1

Met	Arg	Ser	Thr	Ile	Leu	Leu	Phe	Cys	Leu	Leu	Gly	Ser	Thr	Arg
1				5					10					15
Ser	Leu	Pro	Val	Phe	Pro	Ser	Leu	Ser	Leu	Ile	Pro	Leu	Thr	Gln
				20					25					30
Met	Leu	Thr	Leu	Gly	Pro	Asp	Leu	His	Leu	Leu	Asn	Pro	Ala	Ala
				35					40					45
Gly	Met	Thr	Pro	Gly	Thr	Gln	Thr	His	Pro	Leu	Thr	Leu	Gly	Gly
				50					55					60
Leu	Asn	Val	Gln	Gln	Gln	Leu	His	Pro	His	Val	Leu	Pro	Ile	Phe
				65					70					75
Val	Thr	Gln	Leu	Gly	Ala	Pro	Gly	His	Tyr	Pro	Lys	Leu	Arg	Gly
				80					85					90
Ile	Ala	Thr	Asn	Leu	His	Glu	Pro	His	His	Pro	Phe	Leu	Val	Pro
				95					100					105
Arg	Glu	Ala	Ser	Leu	Pro	Thr	Ser	Gln	Ala	Gly	Ala	Asn	Pro	Asp
				110					115					120
Val	Gln	Asp	Gly	Ser	Leu	Pro	Ala	Gly	Gly	Ala	Gly	Val	Asn	Pro
				125					130					135
Ala	Thr	Gln	Gly	Thr	Pro	Ala	Gly	Arg	Leu	Pro	Thr	Pro	Ser	Gly
				140					145					150
Thr	Asp	Asp	Asp	Phe	Ala	Val	Thr	Thr	Pro	Ala	Gly	Ile	Gln	Arg
				155					160					165
Ser	Thr	His	Ala	Ile	Glu	Glu	Ala	Thr	Thr	Glu	Ser	Ala	Asn	Gly
				170					175					180
Ile	Gln													

<210> 2

<211> 125

<212> PRT
 <213> Homo sapiens
 <220> -
 <221> misc_feature
 <223> Incyte ID No: 371156CD1

<400> 2
 Met Val Cys Glu Asp Ala Pro Ser Phe Gln Met Ala Trp Glu Ser
 1 5 10 15
 Gln Met Ala Trp Glu Arg Gly Pro Ala Leu Leu Cys Cys Val Leu
 20 25 30
 Ser Ala Ser Gln Leu Ser Ser Gln Asp Gln Asp Pro Leu Gly His
 35 40 45
 Ile Lys Ser Leu Leu Tyr Pro Phe Gly Phe Pro Val Glu Leu Pro
 50 55 60
 Arg Pro Gly Pro Thr Gly Ala Tyr Lys Lys Val Lys Asn Gln Asn
 65 70 75
 Gln Thr Thr Ser Ser Glu Leu Leu Arg Lys Gln Thr Ser His Phe
 80 85 90
 Asn Gln Arg Gly His Arg Ala Arg Ser Lys Leu Leu Ala Ser Arg
 95 100 105
 Gln Ile Pro Asp Arg Thr Phe Lys Cys Gly Lys Trp Leu Pro Gln
 110 115 120
 Val Pro Ser Pro Val
 125

<210> 3
 <211> 320
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 584050CD1

<400> 3
 Met Ala Gly Leu Ala Ala Arg Leu Val Leu Leu Ala Gly Ala Ala
 1 5 10 15
 Ala Leu Ala Ser Gly Ser Gln Gly Asp Arg Glu Pro Val Tyr Arg
 20 25 30
 Asp Cys Val Leu Gln Cys Glu Glu Gln Asn Cys Ser Gly Gly Ala
 35 40 45
 Leu Asn His Phe Arg Ser Arg Gln Pro Ile Tyr Met Ser Leu Ala
 50 55 60
 Gly Trp Thr Cys Arg Asp Asp Cys Lys Tyr Glu Cys Met Trp Val
 65 70 75
 Thr Val Gly Leu Tyr Leu Gln Glu Gly His Lys Val Pro Gln Phe
 80 85 90
 His Gly Lys Trp Pro Phe Ser Arg Phe Leu Phe Phe Gln Glu Pro
 95 100 105
 Ala Ser Ala Val Ala Ser Phe Leu Asn Gly Leu Ala Ser Leu Val
 110 115 120
 Met Leu Cys Arg Tyr Arg Thr Phe Val Pro Ala Ser Ser Pro Met
 125 130 135
 Tyr His Thr Cys Val Ala Phe Ala Trp Val Ser Leu Asn Ala Trp
 140 145 150
 Phe Trp Ser Thr Val Phe His Thr Arg Asp Thr Asp Leu Thr Glu
 155 160 165
 Lys Met Asp Tyr Phe Cys Ala Ser Thr Val Ile Leu His Ser Ile
 170 175 180
 Tyr Leu Cys Cys Val Arg Thr Val Gly Leu Gln His Pro Ala Val
 185 190 195
 Val Ser Ala Phe Arg Ala Leu Leu Leu Leu Met Leu Thr Val His

Val Ser Tyr Leu	200	Asp Tyr Gly Tyr Asn	210
Ser Leu Ile Arg Phe	205	Leu	215
Val Ala Asn Val	215	Val Val Trp Trp	225
Ala Ile Gly Leu Val	230	Leu	235
Ala Trp Cys Leu	235	Pro His Val Arg	240
Trp Asn Gln Arg Arg	245	Lys	250
Cys Val Val Val	250	Gly Leu Ser Leu Leu	255
Val Leu Leu Leu Gln	260	Glu	265
Leu Leu Asp Phe	265	Val Leu Asp Ala His	270
Pro Pro Leu Phe Trp	275	Ala	280
Ile Trp His Ile	280	His Val Leu Phe Phe	285
Ser Thr Ile Pro Val	290	Ser	295
Phe Leu Glu Asp	295	Leu Lys Glu Ser Glu	300
Asp Ser Leu Tyr Leu	305	Asp	310
Lys Phe Lys Leu Asp	310		315
	320		

<210> 4

<211> 234

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 863808CD1

<400> 4

Met Gly Pro Gly Gly	Arg Val Ala Arg	Leu Leu Ala Pro	Leu Met
1	5	10	15
Trp Arg Arg Ala Val	Ser Ser Val Ala	Gly Ser Ala Val	Gly Ala
	20	25	30
Glu Pro Gly Leu Arg	Leu Leu Ala Val	Gln Arg Leu Pro	Val Gly
	35	40	45
Ala Ala Phe Cys Arg	Ala Cys Gln Thr	Pro Asn Phe Val	Arg Gly
	50	55	60
Leu His Ser Glu Pro	Gly Leu Glu Glu	Arg Ala Glu Gly	Thr Val
	65	70	75
Asn Glu Gly Arg Pro	Glu Ser Asp Ala	Ala Asp His Thr	Gly Pro
	80	85	90
Lys Phe Asp Ile Asp	Met Met Val Ser	Leu Leu Arg Gln	Glu Asn
	95	100	105
Ala Arg Asp Ile Cys	Val Ile Gln Val	Pro Pro Glu Met	Arg Tyr
	110	115	120
Thr Asp Tyr Phe Val	Ile Val Ser Gly	Thr Ser Thr Arg	His Leu
	125	130	135
His Ala Met Ala Phe	Tyr Val Val Lys	Met Tyr Lys His	Leu Lys
	140	145	150
Cys Lys Arg Asp Pro	His Val Lys Ile	Glu Gly Lys Asp	Thr Asp
	155	160	165
Asp Trp Leu Cys Val	Asp Phe Gly Ser	Met Val Ile His	Leu Met
	170	175	180
Leu Pro Glu Thr Arg	Glu Ile Tyr Glu	Leu Glu Lys Leu	Trp Thr
	185	190	195
Leu Arg Ser Tyr Asp	Asp Gln Leu Ala	Gln Ile Ala Pro	Glu Thr
	200	205	210
Val Pro Glu Asp Phe	Ile Leu Gly Ile	Glu Asp Asp Thr	Ser Ser
	215	220	225
Val Thr Pro Val Glu	Leu Lys Cys Glu		
	230		

<210> 5
 <211> 278
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 978433CD1

<400> 5
 Met Gln Pro Ala Ala Ala Ser Glu Arg Gly Gly Ala Asp Ala Asp
 1 5 10 15
 His Val Pro Leu Leu Gly Leu Leu Arg Leu Gln Leu Arg Ala Ala
 20 25 30
 Arg Gln Pro Gly Ala Met Arg Pro Gln Gly Pro Ala Ala Ser Pro
 35 40 45
 Gln Arg Leu Arg Gly Leu Leu Leu Leu Leu Leu Gln Leu Pro
 50 55 60
 Ala Pro Ser Ser Ala Ser Glu Ile Pro Lys Gly Lys Gln Lys Ala
 65 70 75
 Gln Leu Arg Gln Arg Glu Val Val Asp Leu Tyr Asn Gly Met Cys
 80 85 90
 Leu Gln Gly Pro Ala Gly Val Pro Gly Arg Asp Gly Ser Pro Gly
 95 100 105
 Ala Asn Gly Ile Pro Gly Thr Pro Gly Ile Pro Gly Arg Asp Gly
 110 115 120
 Phe Lys Gly Glu Lys Gly Glu Cys Leu Arg Glu Ser Phe Glu Glu
 125 130 135
 Ser Trp Thr Pro Asn Tyr Lys Gln Cys Ser Trp Ser Ser Leu Asn
 140 145 150
 Tyr Gly Ile Asp Leu Gly Lys Ile Ala Glu Cys Thr Phe Thr Lys
 155 160 165
 Met Arg Ser Asn Ser Ala Leu Arg Val Leu Phe Ser Gly Ser Leu
 170 175 180
 Arg Leu Lys Cys Arg Asn Ala Cys Cys Gln Arg Trp Tyr Phe Thr
 185 190 195
 Phe Asn Gly Ala Glu Cys Ser Gly Pro Leu Pro Ile Glu Ala Ile
 200 205 210
 Ile Tyr Leu Asp Gln Gly Ser Pro Glu Met Asn Ser Thr Ile Asn
 215 220 225
 Ile His Arg Thr Ser Ser Val Glu Gly Leu Cys Glu Gly Ile Gly
 230 235 240
 Ala Gly Leu Val Asp Val Ala Ile Trp Val Gly Thr Cys Ser Asp
 245 250 255
 Tyr Pro Lys Gly Asp Ala Ser Thr Gly Trp Asn Ser Val Ser Arg
 260 265 270
 Ile Ile Ile Glu Glu Leu Pro Lys
 275

<210> 6
 <211> 136
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1655369CD1

<400> 6
 Met Pro Pro Gly Gly Leu Gly Ala Cys Ala Val Thr Pro Ala Pro
 1 5 10 15
 Gly Glu Glu Arg Thr Gln Pro Gly Glu Leu Gly Gln Gly Leu His
 20 25 30
 Met Ala Gln Gly Gln Gln Met Leu Ala Gly Gln Leu Leu Pro Met

				35					40					45
Leu	Thr	Leu	Leu	Pro	Pro	Ser	Phe	Pro	Leu	Pro	His	Pro	Thr	Leu
				50					55					60
Gly	Pro	Arg	Arg	His	Ala	Ser	Leu	Thr	Gln	Leu	Gly	Pro	Ala	Phe
				65					70					75
Trp	Met	Ala	Trp	Gly	Arg	Pro	Trp	Ala	His	Leu	Gly	Pro	Gly	Gln
				80					85					90
Pro	Leu	Gly	Gln	Leu	Trp	Lys	Ser	Ser	Val	Glu	Glu	His	Leu	Leu
				95					100					105
Ala	Ala	Trp	Leu	Gln	Pro	Leu	Ala	Leu	Leu	Glu	Trp	Ser	Leu	Gly
				110					115					120
Ala	Ser	Ala	Leu	Ser	Ala	Leu	Gly	Thr	Ser	His	Pro	Leu	Gly	Leu
				125					130					135
Gln														

<210> 7

<211> 109

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1703244CD1

<400> 7

Met	Leu	Met	Tyr	Met	Phe	Tyr	Val	Leu	Pro	Phe	Cys	Gly	Leu	Ala
1				5					10					15
Ala	Tyr	Ala	Leu	Thr	Phe	Pro	Gly	Cys	Ser	Trp	Leu	Pro	Asp	Trp
				20					25					30
Ala	Leu	Val	Phe	Ala	Gly	Gly	Ile	Gly	Gln	Ala	Gln	Phe	Ser	His
				35					40					45
Met	Gly	Ala	Ser	Met	His	Leu	Arg	Thr	Pro	Phe	Thr	Tyr	Arg	Val
				50					55					60
Pro	Glu	Asp	Thr	Trp	Gly	Cys	Phe	Phe	Val	Cys	Asn	Leu	Leu	Tyr
				65					70					75
Ala	Leu	Gly	Pro	His	Leu	Leu	Ala	Tyr	Arg	Cys	Leu	Gln	Trp	Pro
				80					85					90
Ala	Phe	Phe	His	Gln	Pro	Pro	Pro	Ser	Asp	Pro	Leu	Ala	Leu	His
				95					100					105
Lys	Lys	Gln	His											

<210> 8

<211> 262

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1730819CD1

<400> 8

Met	Ala	Ala	Ala	Ser	Ala	Gly	Ala	Thr	Arg	Leu	Leu	Leu	Leu	Leu
1				5					10					15
Leu	Met	Ala	Val	Ala	Ala	Pro	Ser	Arg	Ala	Arg	Gly	Ser	Gly	Cys
				20					25					30
Arg	Ala	Gly	Thr	Gly	Ala	Arg	Gly	Ala	Gly	Ala	Glu	Gly	Arg	Glu
				35					40					45
Gly	Glu	Ala	Cys	Gly	Thr	Val	Gly	Leu	Leu	Leu	Glu	His	Ser	Phe
				50					55					60
Glu	Ile	Asp	Asp	Ser	Ala	Asn	Phe	Arg	Lys	Arg	Gly	Ser	Leu	Leu
				65					70					75

```

Trp Asn Gln Gln Asp Gly Thr Leu Ser Leu Ser Gln Arg Gln Leu
      80      85      90
Ser Glu Glu Glu Arg Gly Arg Leu Arg Asp Val Ala Ala Leu Asn
      95     100     105
Gly Leu Tyr Arg Val Arg Ile Pro Arg Arg Pro Gly Ala Leu Asp
     110     115     120
Gly Leu Glu Ala Gly Gly Tyr Val Ser Ser Phe Val Pro Ala Cys
     125     130     135
Ser Leu Val Glu Ser His Leu Ser Asp Gln Leu Thr Leu His Val
     140     145     150
Asp Val Ala Gly Asn Val Val Gly Val Ser Val Val Thr His Pro
     155     160     165
Gly Gly Cys Arg Gly His Glu Val Glu Asp Val Asp Leu Glu Leu
     170     175     180
Phe Asn Thr Ser Val Gln Leu Gln Pro Pro Thr Thr Ala Pro Gly
     185     190     195
Pro Glu Thr Ala Ala Phe Ile Glu Arg Leu Glu Met Glu Gln Ala
     200     205     210
Gln Lys Ala Lys Asn Pro Gln Glu Gln Lys Ser Phe Phe Ala Lys
     215     220     225
Tyr Trp Met Tyr Ile Ile Pro Val Val Leu Phe Leu Met Met Ser
     230     235     240
Gly Ala Pro Asp Thr Gly Gly Gln Gly Gly Gly Gly Gly Cys Gly
     245     250     255
Gly Gly Gly Gly Ser Gly Arg
      260

```

<210> 9

<211> 384

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1757161CD1

<400> 9

```

Met Ala Glu Gln Thr Tyr Ser Trp Ala Tyr Ser Leu Val Asp Ser
  1      5      10      15
Ser Gln Val Ser Thr Phe Leu Ile Ser Ile Leu Leu Ile Val Tyr
      20      25      30
Gly Ser Phe Arg Ser Leu Asn Met Asp Phe Glu Asn Gln Asp Lys
      35      40      45
Glu Lys Asp Ser Asn Ser Ser Ser Gly Ser Phe Asn Gly Asn Ser
      50      55      60
Thr Asn Asn Ser Ile Gln Thr Ile Asp Ser Thr Gln Ala Leu Phe
      65      70      75
Leu Pro Ile Gly Ala Ser Val Ser Leu Leu Val Met Phe Phe Phe
      80      85      90
Phe Asp Ser Val Gln Val Val Phe Thr Ile Cys Thr Ala Val Leu
      95     100     105
Ala Thr Ile Ala Phe Ala Phe Leu Leu Leu Pro Met Cys Gln Tyr
     110     115     120
Leu Thr Arg Pro Cys Ser Pro Gln Asn Lys Ile Ser Phe Gly Cys
     125     130     135
Cys Gly Arg Phe Thr Ala Ala Glu Leu Leu Ser Phe Ser Leu Ser
     140     145     150
Val Met Leu Val Leu Ile Trp Val Leu Thr Gly His Trp Leu Leu
     155     160     165
Met Asp Ala Leu Ala Met Gly Leu Cys Val Ala Met Ile Ala Phe
     170     175     180
Val Arg Leu Pro Ser Leu Lys Val Ser Cys Leu Leu Leu Ser Gly
     185     190     195
Leu Leu Ile Tyr Asp Val Phe Trp Val Phe Phe Ser Ala Tyr Ile

```

Phe Asn Ser Asn Val Met Val Lys Val	200	Ala Thr Gln Pro Ala Asp	205	210
Asn Pro Leu Asp Val Leu Ser Arg Lys	215	Leu His Leu Gly Pro Asn	220	225
Val Gly Arg Asp Val Pro Arg Leu Ser	230	Leu Pro Gly Lys Leu Val	235	240
Phe Pro Ser Ser Thr Gly Ser His Phe	245	Ser Met Leu Gly Ile Gly	250	255
Asp Ile Val Met Pro Gly Leu Leu Leu	260	Cys Phe Val Leu Arg Tyr	265	270
Asp Asn Tyr Lys Lys Gln Ala Ser Gly	275	Ser Cys Gly Ala Pro	280	285
Gly Pro Ala Asn Ile Ser Gly Arg Met	290	Gln Lys Val Ser Tyr Phe	295	300
His Cys Thr Leu Ile Gly Tyr Phe Val	305	Gly Leu Leu Thr Ala Thr	310	315
Val Ala Ser Arg Ile His Arg Ala Ala	320	Gln Pro Ala Leu Leu Tyr	325	330
Leu Val Pro Phe Thr Leu Leu Pro Leu	335	Leu Thr Met Ala Tyr Leu	340	345
Lys Gly Asp Leu Arg Arg Met Trp Ser	350	Glu Pro Phe His Ser Lys	355	360
Ser Ser Ser Ser Arg Phe Leu Glu Val	365		370	375
	380			

<210> 10
 <211> 244
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1976095CD1

Met Asp Ile Leu Val Pro Leu Leu Gln Leu Leu Val Leu Leu Leu	<400> 10	
1 5 10 15		
Thr Leu Pro Leu His Leu Met Ala Leu Leu Gly Cys Trp Gln Pro	20 25 30	
Leu Cys Lys Ser Tyr Phe Pro Tyr Leu Met Ala Val Leu Thr Pro	35 40 45	
Lys Ser Asn Arg Lys Met Glu Ser Lys Lys Arg Glu Leu Phe Ser	50 55 60	
Gln Ile Lys Gly Leu Thr Gly Ala Ser Gly Lys Val Ala Leu Leu	65 70 75	
Glu Leu Gly Cys Gly Thr Gly Ala Asn Phe Gln Phe Tyr Pro Pro	80 85 90	
Gly Cys Arg Val Thr Cys Leu Asp Pro Asn Pro His Phe Glu Lys	95 100 105	
Phe Leu Thr Lys Ser Met Ala Glu Asn Arg His Leu Gln Tyr Glu	110 115 120	
Arg Phe Val Val Ala Pro Gly Glu Asp Met Arg Gln Leu Ala Asp	125 130 135	
Gly Ser Met Asp Val Val Val Cys Thr Leu Val Leu Cys Ser Val	140 145 150	
Gln Ser Pro Arg Lys Val Leu Gln Glu Val Arg Arg Val Leu Arg	155 160 165	
Pro Gly Gly Val Leu Phe Phe Trp Glu His Val Ala Glu Pro Tyr	170 175 180	
Gly Ser Trp Ala Phe Met Trp Gln Gln Val Phe Glu Pro Thr Trp	185 190 195	
Lys His Ile Gly Asp Gly Cys Cys Leu Thr Arg Glu Thr Trp Lys	200 205 210	

Asp	Leu	Glu	Asn	Ala	Gln	Phe	Ser	Glu	Ile	Gln	Met	Glu	Arg	Gln
				215					220					225
Pro	Pro	Pro	Leu	Lys	Trp	Leu	Pro	Val	Gly	Pro	His	Ile	Met	Gly
				230					235					240
Lys	Ala	Val	Lys											

<210> 11
 <211> 326
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2169991CD1

<400> 11

Met	Arg	Thr	Glu	Ala	Gln	Val	Pro	Ala	Leu	Gln	Pro	Pro	Glu	Pro
1				5					10					15
Gly	Leu	Glu	Gly	Ala	Met	Gly	His	Arg	Thr	Leu	Val	Leu	Pro	Trp
				20					25					30
Val	Leu	Leu	Thr	Leu	Cys	Val	Thr	Ala	Gly	Thr	Pro	Glu	Val	Trp
				35					40					45
Val	Gln	Val	Arg	Met	Glu	Ala	Thr	Glu	Leu	Ser	Ser	Phe	Thr	Ile
				50					55					60
Arg	Cys	Gly	Phe	Leu	Gly	Ser	Gly	Ser	Ile	Ser	Leu	Val	Thr	Val
				65					70					75
Ser	Trp	Gly	Gly	Pro	Asn	Gly	Ala	Gly	Gly	Thr	Thr	Leu	Ala	Val
				80					85					90
Leu	His	Pro	Glu	Arg	Gly	Ile	Arg	Gln	Trp	Ala	Pro	Ala	Arg	Gln
				95					100					105
Ala	Arg	Trp	Glu	Thr	Gln	Ser	Ser	Ile	Ser	Leu	Ile	Leu	Glu	Gly
				110					115					120
Ser	Gly	Ala	Ser	Ser	Pro	Cys	Ala	Asn	Thr	Thr	Phe	Cys	Cys	Lys
				125					130					135
Phe	Ala	Ser	Phe	Pro	Glu	Gly	Ser	Trp	Glu	Ala	Cys	Gly	Ser	Leu
				140					145					150
Pro	Pro	Ser	Ser	Asp	Pro	Gly	Leu	Ser	Ala	Pro	Pro	Thr	Pro	Ala
				155					160					165
Pro	Ile	Leu	Arg	Ala	Asp	Leu	Ala	Gly	Ile	Leu	Gly	Val	Ser	Gly
				170					175					180
Val	Leu	Leu	Phe	Gly	Cys	Val	Tyr	Leu	Leu	His	Leu	Leu	Arg	Arg
				185					190					195
His	Lys	His	Arg	Pro	Ala	Pro	Arg	Leu	Gln	Pro	Ser	Arg	Thr	Ser
				200					205					210
Pro	Gln	Ala	Pro	Arg	Ala	Arg	Ala	Trp	Ala	Pro	Ser	Gln	Ala	Ser
				215					220					225
Gln	Ala	Ala	Leu	His	Val	Pro	Tyr	Ala	Thr	Ile	Asn	Thr	Ser	Cys
				230					235					240
Arg	Pro	Ala	Thr	Leu	Asp	Thr	Ala	His	Pro	His	Gly	Gly	Pro	Ser
				245					250					255
Trp	Trp	Ala	Ser	Leu	Pro	Thr	His	Ala	Ala	His	Arg	Pro	Gln	Gly
				260					265					270
Pro	Ala	Ala	Trp	Ala	Ser	Thr	Pro	Ile	Pro	Ala	Arg	Gly	Ser	Phe
				275					280					285
Val	Ser	Val	Glu	Asn	Gly	Leu	Tyr	Ala	Gln	Ala	Gly	Glu	Arg	Pro
				290					295					300
Pro	His	Thr	Gly	Pro	Gly	Leu	Thr	Leu	Phe	Pro	Asp	Pro	Arg	Gly
				305					310					315
Pro	Arg	Ala	Met	Glu	Gly	Pro	Leu	Gly	Val	Arg				
				320					325					

<210> 12
 <211> 105
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2616827CD1

<400> 12
 Met Asn Leu Gly Val Ser Met Leu Arg Ile Leu Phe Leu Leu Asp
 1 5 10 15
 Val Gly Gly Ala Gln Val Leu Ala Thr Gly Lys Thr Pro Gly Ala
 20 25 30
 Glu Ile Asp Phe Lys Tyr Ala Leu Ile Gly Thr Ala Val Gly Val
 35 40 45
 Ala Ile Ser Ala Gly Phe Leu Ala Leu Lys Ile Cys Met Ile Arg
 50 55 60
 Arg His Leu Phe Asp Asp Ser Ser Asp Leu Lys Ser Thr Pro
 65 70 75
 Gly Gly Leu Ser Asp Thr Ile Pro Leu Lys Lys Arg Ala Pro Arg
 80 85 90
 Arg Asn His Asn Phe Ser Lys Arg Asp Ala Gln Val Ile Glu Leu
 95 100 105

<210> 13
 <211> 626
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2991370CD1

<400> 13
 Met Ala Pro Ser Ala Asp Pro Gly Met Ser Arg Met Leu Pro Phe
 1 5 10 15
 Leu Leu Leu Leu Trp Phe Leu Pro Ile Thr Glu Gly Ser Gln Arg
 20 25 30
 Ala Glu Pro Met Phe Thr Ala Val Thr Asn Ser Val Leu Pro Pro
 35 40 45
 Asp Tyr Asp Ser Asn Pro Thr Gln Leu Asn Tyr Gly Val Ala Val
 50 55 60
 Thr Asp Val Asp His Asp Gly Asp Phe Glu Ile Val Val Ala Gly
 65 70 75
 Tyr Asn Gly Pro Asn Leu Val Leu Lys Tyr Asp Arg Ala Gln Lys
 80 85 90
 Arg Leu Val Asn Ile Ala Val Asp Glu Arg Ser Ser Pro Tyr Tyr
 95 100 105
 Ala Leu Arg Asp Arg Gln Gly Asn Ala Ile Gly Val Thr Ala Cys
 110 115 120
 Asp Ile Asp Gly Asp Gly Arg Glu Glu Ile Tyr Phe Leu Asn Thr
 125 130 135
 Asn Asn Ala Phe Ser Gly Val Ala Thr Tyr Thr Asp Lys Leu Phe
 140 145 150
 Lys Phe Arg Asn Asn Arg Trp Glu Asp Ile Leu Ser Asp Glu Val
 155 160 165
 Asn Val Ala Arg Gly Val Ala Ser Leu Phe Ala Gly Arg Ser Val
 170 175 180
 Ala Cys Val Asp Arg Lys Gly Ser Gly Arg Tyr Ser Ile Tyr Ile
 185 190 195
 Ala Asn Tyr Ala Tyr Gly Asn Val Gly Pro Asp Ala Leu Ile Glu
 200 205 210
 Met Asp Pro Glu Ala Ser Asp Leu Ser Arg Gly Ile Leu Ala Leu

Arg Asp Val Ala	215	Ala Glu Ala Gly Val	220	Ser Lys Tyr Thr Gly	225
Arg Gly Val Ser	230	Val Gly Pro Ile Leu	235	Ser Ser Ser Ala Ser	240
Ile Phe Cys Asp	245	Asn Glu Asn Gly Pro	250	Asn Phe Leu Phe His	255
Arg Gly Asp Gly	260	Thr Phe Val Asp Ala	265	Ala Ala Ser Ala Gly	270
Asp Asp Pro His	275	Gln His Gly Arg Gly	280	Val Ala Leu Ala Asp	285
Asn Arg Asp Gly	290	Lys Val Asp Ile Val	295	Tyr Gly Asn Trp Asn	300
Pro His Arg Leu	305	Tyr Leu Gln Met Ser	310	Thr His Gly Lys Val	315
Phe Arg Asp Ile	320	Ala Ser Pro Lys Phe	325	Ser Met Pro Ser Pro	330
Arg Thr Val Ile	335	Thr Ala Asp Phe Asp	340	Asn Asp Gln Glu Leu	345
Ile Phe Phe Asn	350	Asn Ile Ala Tyr Arg	355	Ser Ser Ser Ala Asn	360
Leu Phe Arg Val	365	Ile Arg Arg Glu His	370	Gly Asp Pro Leu Ile	375
Glu Leu Asn Pro	380	Gly Asp Ala Leu Glu	385	Pro Glu Gly Arg Gly	390
Gly Gly Val Val	395	Thr Asp Phe Asp Gly	400	Asp Gly Met Leu Asp	405
Ile Leu Ser His	410	Gly Glu Ser Met Ala	415	Gln Pro Leu Ser Val	420
Arg Gly Asn Gln	425	Gly Phe Asn Asn Asn	430	Trp Leu Arg Val Val	435
Arg Thr Arg Phe	440	Gly Ala Phe Ala Arg	445	Gly Ala Lys Val Val	450
Tyr Thr Lys Lys	455	Ser Gly Ala His Leu	460	Arg Ile Ile Asp Gly	465
Ser Gly Tyr Leu	470	Cys Glu Met Glu Pro	475	Val Ala His Phe Gly	480
Gly Lys Asp Glu	485	Ala Ser Ser Val Glu	490	Val Thr Trp Pro Asp	495
Lys Met Val Ser	500	Arg Asn Val Ala Ser	505	Gly Glu Met Asn Ser	510
Leu Glu Ile Leu	515	Tyr Pro Arg Asp Glu	520	Asp Thr Leu Gln Asp	525
Ala Pro Leu Glu	530	Cys Gly Gln Gly Phe	535	Ser Gln Gln Glu Asn	540
His Cys Met Asp	545	Thr Asn Glu Cys Ile	550	Gln Phe Pro Phe Val	555
Pro Arg Asp Lys	560	Pro Val Cys Val Asn	565	Thr Tyr Gly Ser Tyr	570
Cys Arg Thr Asn	575	Lys Lys Cys Ser Arg	580	Gly Tyr Glu Pro Asn	585
Asp Gly Thr Ala	590	Cys Val Gly Trp Trp	595	Ser Pro Val Leu Lys	600
Val Thr Pro Gln	605	Val Gly Lys Ser Leu	610	Gly Pro	615
	620		625		

<210> 14

<211> 296

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3031062CD1

<400> 14

```

Met Glu Trp Trp Ala Ser Ser Pro Leu Arg Leu Trp Leu Leu Leu
 1      5      10      15
Phe Leu Leu Pro Ser Ala Gln Gly Arg Gln Lys Glu Ser Gly Ser
 20      25      30
Lys Trp Lys Val Phe Ile Asp Gln Ile Asn Arg Ser Leu Glu Asn
 35      40      45
Tyr Glu Pro Cys Ser Ser Gln Asn Cys Ser Cys Tyr His Gly Val
 50      55      60
Ile Glu Glu Asp Leu Thr Pro Phe Arg Gly Gly Ile Ser Arg Lys
 65      70      75
Met Met Ala Glu Val Val Arg Arg Lys Leu Gly Thr His Tyr Gln
 80      85      90
Ile Thr Lys Asn Arg Leu Tyr Arg Glu Asn Asp Cys Met Phe Pro
 95      100      105
Ser Arg Cys Ser Gly Val Glu His Phe Ile Leu Glu Val Ile Gly
 110      115      120
Arg Leu Pro Asp Met Glu Met Val Ile Asn Val Arg Asp Tyr Pro
 125      130      135
Gln Val Pro Lys Trp Met Glu Pro Ala Ile Pro Val Phe Ser Phe
 140      145      150
Ser Lys Thr Ser Glu Tyr His Asp Ile Met Tyr Pro Ala Trp Thr
 155      160      165
Phe Trp Glu Gly Gly Pro Ala Val Trp Pro Ile Tyr Pro Thr Gly
 170      175      180
Leu Gly Arg Trp Asp Leu Phe Arg Glu Asp Leu Val Arg Ser Ala
 185      190      195
Ala Gln Trp Pro Trp Lys Lys Lys Asn Ser Thr Ala Tyr Phe Arg
 200      205      210
Gly Ser Arg Thr Ser Pro Glu Arg Asp Pro Leu Ile Leu Leu Ser
 215      220      225
Arg Lys Asn Pro Lys Leu Val Asp Ala Glu Tyr Thr Lys Asn Gln
 230      235      240
Ala Trp Lys Ser Met Lys Asp Thr Leu Gly Lys Pro Ala Ala Lys
 245      250      255
Asp Val His Leu Val Asp His Cys Lys Tyr Lys Tyr Leu Phe Asn
 260      265      270
Phe Arg Gly Val Leu Gln Val Ser Gly Leu Asn Thr Ser Ser Cys
 275      280      285
Val Ala Ile Ile Leu Met Arg Lys Arg Thr Tyr
 290      295

```

<210> 15

<211> 249

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3101617CD1

<400> 15

```

Met Asp Gly Lys Lys Cys Ser Val Trp Met Phe Leu Pro Leu Val
 1      5      10      15
Phe Thr Leu Phe Thr Ser Ala Gly Leu Trp Ile Val Tyr Phe Ile
 20      25      30
Ala Val Glu Asp Asp Lys Ile Leu Pro Leu Asn Ser Ala Glu Arg
 35      40      45
Lys Pro Gly Val Lys His Ala Pro Tyr Ile Ser Ile Ala Gly Asp
 50      55      60
Asp Pro Pro Ala Ser Cys Val Phe Ser Gln Val Met Asn Met Ala
 65      70      75
Ala Phe Leu Ala Leu Val Val Ala Val Leu Arg Phe Ile Gln Leu
 80      85      90

```

```

Lys Pro Lys Val Leu Asn Pro Trp Leu Asn Ile Ser Gly Leu Val
      95      100      105
Ala Leu Cys Leu Ala Ser Phe Gly Met Thr Leu Leu Gly Asn Phe
      110      115      120
Gln Leu Thr Asn Asp Glu Glu Ile His Asn Val Gly Thr Ser Leu
      125      130      135
Thr Phe Gly Phe Gly Thr Leu Thr Cys Trp Ile Gln Ala Ala Leu
      140      145      150
Thr Leu Lys Val Asn Ile Lys Asn Glu Gly Arg Arg Val Gly Ile
      155      160      165
Pro Arg Val Ile Leu Ser Ala Ser Ile Thr Leu Cys Val Val Leu
      170      175      180
Tyr Phe Ile Leu Met Ala Gln Ser Ile His Met Tyr Ala Ala Arg
      185      190      195
Val Gln Trp Gly Leu Val Met Cys Phe Leu Ser Tyr Phe Gly Thr
      200      205      210
Phe Ala Val Glu Phe Arg His Tyr Arg Tyr Glu Ile Val Cys Ser
      215      220      225
Glu Tyr Gln Glu Asn Phe Leu Ser Phe Ser Glu Ser Leu Ser Glu
      230      235      240
Ala Ser Glu Tyr Gln Thr Asp Gln Val
      245

```

<210> 16
 <211> 124
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3216178CD1

```

<400> 16
Met Gly Gly Tyr Leu Lys Thr Arg Pro Trp Thr Leu Gln His Phe
  1      5      10      15
Tyr Leu Cys Leu Met Pro Ala Ala Thr Trp Leu Val Leu Leu Leu
      20      25      30
Leu Leu Trp Leu Ser Leu Gly Val Lys Thr Gly Ser Cys Ser Gln
      35      40      45
Pro Gln Asn Leu Cys Cys Leu Gly Thr Asp His His Cys Lys Arg
      50      55      60
Gly Ser Cys Tyr Cys Asp Glu Phe Cys His Val Ala Pro Asp Cys
      65      70      75
His Pro Asp His Ser Val Leu Cys Asn Pro Ala Ser Gln Met Thr
      80      85      90
Lys Met Val Leu Gln Met Val Leu Arg Met Glu Asn Pro Pro Ser
      95      100      105
Pro Ala Arg Ser His Leu Asp Trp Met Gln Ser Met Val Ser Ser
      110      115      120
Leu Gln Val Leu

```

<210> 17
 <211> 101
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3406803CD1

<400> 17

Met	Leu	Pro	Val	Gly	Ala	Gln	Pro	Arg	Ser	Pro	Pro	Trp	Val	Leu
1				5					10					15
Ala	Arg	Leu	Leu	His	Pro	Arg	Gly	Pro	Ala	Ala	Thr	Ser	Leu	Val
				20					25					30
Pro	Phe	Leu	Pro	Trp	Gly	Ser	Leu	Glu	Ser	His	Thr	Pro	Cys	Pro
				35					40					45
Tyr	Arg	Ala	Cys	Ser	Pro	Gly	Trp	Glu	Leu	Thr	Leu	Ser	Thr	Phe
				50					55					60
Pro	Glu	Arg	Glu	Thr	Leu	Ser	Gly	Gly	Glu	Val	Arg	Lys	Arg	Gly
				65					70					75
Ala	Gly	Ser	Met	Val	Gly	Gly	Gly	Glu	Ser	Thr	Met	Thr	Arg	Ala
				80					85					90
Leu	Cys	Val	Arg	Leu	Leu	Thr	Lys	Leu	Arg	Val				
				95					100					

<210> 18

<211> 540

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3468066CD1

<400> 18

Met	Ala	Thr	Ser	Gly	Ala	Ala	Ser	Ala	Glu	Leu	Val	Ile	Gly	Trp
1				5					10					15
Cys	Ile	Phe	Gly	Leu	Leu	Leu	Leu	Ala	Ile	Leu	Ala	Phe	Cys	Trp
				20					25					30
Ile	Tyr	Val	Arg	Lys	Tyr	Gln	Ser	Arg	Arg	Glu	Ser	Glu	Val	Val
				35					40					45
Ser	Thr	Ile	Thr	Ala	Ile	Phe	Ser	Leu	Ala	Ile	Ala	Leu	Ile	Thr
				50					55					60
Ser	Ala	Leu	Leu	Pro	Val	Asp	Ile	Phe	Leu	Val	Ser	Tyr	Met	Lys
				65					70					75
Asn	Gln	Asn	Gly	Thr	Phe	Lys	Asp	Trp	Ala	Asn	Ala	Asn	Val	Ser
				80					85					90
Arg	Gln	Ile	Glu	Asp	Thr	Val	Leu	Tyr	Gly	Tyr	Tyr	Thr	Leu	Tyr
				95					100					105
Ser	Val	Ile	Leu	Phe	Cys	Val	Phe	Phe	Trp	Ile	Pro	Phe	Val	Tyr
				110					115					120
Phe	Tyr	Tyr	Glu	Glu	Lys	Asp	Asp	Asp	Asp	Thr	Ser	Lys	Cys	Thr
				125					130					135
Gln	Ile	Lys	Thr	Ala	Leu	Lys	Tyr	Thr	Leu	Gly	Phe	Val	Val	Ile
				140					145					150
Cys	Ala	Leu	Leu	Leu	Leu	Val	Gly	Ala	Phe	Val	Pro	Leu	Asn	Val
				155					160					165
Pro	Asn	Asn	Lys	Asn	Ser	Thr	Glu	Trp	Glu	Lys	Val	Lys	Ser	Leu
				170					175					180
Phe	Glu	Glu	Leu	Gly	Ser	Ser	His	Gly	Leu	Ala	Ala	Leu	Ser	Phe
				185					190					195
Ser	Ile	Ser	Ser	Leu	Thr	Leu	Ile	Gly	Met	Leu	Ala	Ala	Ile	Thr
				200					205					210
Tyr	Thr	Ala	Tyr	Gly	Met	Ser	Ala	Leu	Pro	Leu	Asn	Leu	Ile	Lys
				215					220					225
Gly	Thr	Arg	Ser	Ala	Ala	Tyr	Glu	Arg	Leu	Glu	Asn	Thr	Glu	Asp
				230					235					240
Ile	Glu	Glu	Val	Glu	Gln	His	Ile	Gln	Thr	Ile	Lys	Ser	Lys	Ser
				245					250					255
Lys	Asp	Gly	Arg	Pro	Leu	Pro	Ala	Arg	Asp	Lys	Arg	Ala	Leu	Lys
				260					265					270
Gln	Phe	Glu	Glu	Arg	Leu	Arg	Thr	Leu	Lys	Lys	Arg	Glu	Arg	His
				275					280					285
Leu	Glu	Phe	Ile	Glu	Asn	Ser	Trp	Trp	Thr	Lys	Phe	Cys	Gly	Ala

14/28

<211> 114
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3669422CD1

<400> 20
 Met Ser Ser Ser Ser Ser Arg Cys Leu Ser Pro Ser Pro Gly Met
 1 5 10 15
 Ser Leu Trp Ser Cys Leu Leu Phe Leu Cys Thr Pro Ser Pro Thr
 20 25 30
 Thr Thr Ser Pro Ser Pro Asp Pro Ser Gln Val Ser Thr Leu Pro
 35 40 45
 Thr Pro Ser Pro Gln Arg Glu Gly Leu Lys Gln Gly Gln Trp Arg
 50 55 60
 Lys Thr Gly Pro Ser Ser Thr His Pro His Thr Pro Ser Ser Arg
 65 70 75
 Pro Pro Ser Pro Ser Ser Leu Pro Leu Thr Trp Lys Leu Leu Gln
 80 85 90
 Pro Ile Pro Ser His Ser Leu Pro His Pro Pro Lys Ile His Thr
 95 100 105
 Gly Pro Ser Leu Ala Glu Cys Gly His
 110

<210> 21
 <211> 114
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3688740CD1

<400> 21
 Met Arg Gly Glu His Asn Ser Thr Ser Tyr Asp Ser Ala Val Ile
 1 5 10 15
 Tyr Arg Gly Phe Trp Ala Val Leu Met Leu Leu Gly Val Val Ala
 20 25 30
 Val Val Ile Ala Ser Phe Leu Ile Ile Cys Ala Ala Pro Phe Ala
 35 40 45
 Ser His Phe Leu Tyr Lys Ala Gly Gly Gly Ser Tyr Ile Ala Ala
 50 55 60
 Asp Gly Ile Ser Ser Leu Cys Tyr Ser Ser Leu Ser Lys Ser Leu
 65 70 75
 Leu Ser Gln Pro Leu Arg Glu Thr Ser Ser Ala Ile Asn Asp Ile
 80 85 90
 Ser Leu Leu Gln Ala Leu Met Pro Leu Leu Gly Trp Thr Ser His
 95 100 105
 Trp Thr Cys Ile Thr Val Gly Leu Tyr
 110

<210> 22
 <211> 287
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3742589CD1

<400> 22

```

Met Glu Leu Glu Arg Ile Val Ser Ala Ala Leu Leu Ala Phe Val
  1          5          10          15
Gln Thr His Leu Pro Glu Ala Asp Leu Ser Gly Leu Asp Glu Val
          20          25          30
Ile Phe Ser Tyr Val Leu Gly Val Leu Glu Asp Leu Gly Pro Ser
          35          40          45
Gly Pro Ser Glu Glu Asn Phe Asp Met Glu Ala Phe Thr Glu Met
          50          55          60
Met Glu Ala Tyr Val Pro Gly Phe Ala His Ile Pro Arg Gly Thr
          65          70          75
Ile Gly Asp Met Met Gln Lys Leu Ser Gly Gln Leu Ser Asp Ala
          80          85          90
Arg Asn Lys Glu Asn Leu Gln Pro Gln Ser Ser Gly Val Gln Gly
          95          100          105
Gln Val Pro Ile Ser Pro Glu Pro Leu Gln Arg Pro Glu Met Leu
          110          115          120
Lys Glu Glu Thr Arg Ser Ser Ala Ala Ala Ala Asp Thr Gln
          125          130          135
Asp Glu Ala Thr Gly Ala Glu Glu Glu Leu Leu Pro Gly Val Asp
          140          145          150
Val Leu Leu Glu Val Phe Pro Thr Cys Ser Val Glu Gln Ala Gln
          155          160          165
Trp Val Leu Ala Lys Ala Arg Gly Asp Leu Glu Glu Ala Val Gln
          170          175          180
Met Leu Val Glu Gly Lys Glu Glu Gly Pro Ala Ala Trp Glu Gly
          185          190          195
Pro Asn Gln Asp Leu Pro Arg Arg Leu Arg Gly Pro Gln Lys Asp
          200          205          210
Glu Leu Lys Ser Phe Ile Leu Gln Lys Tyr Met Met Val Asp Ser
          215          220          225
Ala Glu Asp Gln Lys Ile His Arg Pro Met Ala Pro Lys Glu Ala
          230          235          240
Pro Lys Lys Leu Ile Arg Tyr Ile Asp Asn Gln Val Val Ser Thr
          245          250          255
Lys Gly Glu Arg Phe Lys Asp Val Arg Asn Pro Glu Ala Glu Glu
          260          265          270
Met Lys Ala Thr Tyr Ile Asn Leu Lys Pro Ala Arg Lys Tyr Arg
          275          280          285
Phe His

```

<210> 23

<211> 854

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 078811CB1

<400> 23

```

attttgccctc gtggacccaa aggtagcaat ttgaaacatg aggagtacga ttctactggt 60
ttgtcttcta ggatcaactc gggtcattacc agtctttcct tctttaagtc tgataccatt 120
aacacagatg ctacactgg ggccagatct gcatctgtta aatcctgctg caggaatgac 180
acctggtacc cagaccacc cattgacctt acttggagcc ccagggcact atcctaagct 240
cccacatgtg ttaccaattt ttgtcacaca acttggagcc ccagggcact atcctaagct 300
cagaggaatt gccacaaatc ttcacgagcc tcatcatcca ttccttggtc cccgggaggc 360
atccttgccc accagtcagg caggggctaa tccagatgtc caggatggaa gccttccagc 420
aggaggagca ggtgtaaata ctgccaccca gggaacccca gcaggccgcc tcccaactcc 480
cagtggcaca gatgacgact ttgcagtgc caccctgca ggcattccaa ggagcacaca 540
tgccatcgag gaagccacca cagaatcagc aaatggaatt cagtaagctg tttcaaattt 600
tttcaactaa gctgcctcga atttggtgat acatgtgaat ctttatcatt gattatatta 660
tggaatagat tgagacacat tggatagtct tagaagaaat taattcttaa tttacctgaa 720
aatattcctg aaatttcaga aaatatgttc tatgtagaga atcccaactt ttaaaaaaca 780

```

taattcaatg gataaatctg tctttgaaat ataacattat gctgcctgga tgatatgcat 840
 attaaaacga atta 854

<210> 24
 <211> 1804
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 371156CB1

<400> 24
 gtgataggca gctttccttc ttttcaacag tgatacctac gaaaatcaaa ataaatgcaa 60
 gctgagggtt tgtgctcact gaaagggtcg tcaacccag aaggccgaca caaaaaaat 120
 ggtatgtgaa gatgcaccgt cttttcaaag ggcctgggag agtcaaatgg cctgggagag 180
 ggggcctgcc cttctctgct gtgtcctttc ggcttcccag ttgagctccc aagaccagga 240
 cccactgggg catataaaat ctctgctgta tcccttcggc tccccagttg agtcccaag 300
 accaggaccc actggggcat ataaaaaagt caaaaatcaa aatcaaacia caagttctga 360
 gttacttagg aaacagactt cgcatttcaa tcagagaggg cacagagcaa ggtctaaact 420
 tctggcttct agacaaattc ctgatagaac atttaaagt gggaaagtgg tccccaggt 480
 cccatccccct gtttagggat agagttgata tcatTTTTat aggtgccatg tatgcctctg 540
 cctgaattttt ttttaattgac ttttgagctt ttgagattgc acgagggaga acaaggcctt 600
 tgctgttggtg gataggaaaag acttaaaccta aaatttaaacc agcaagaaaag cattagtaaa 660
 aatctaacia tatgaagggc tcttatgagt catttttttc aaaagatgaa aactccagaa 720
 acgcacagga acgaaatacc tcccagaac atgaagcaat catcgaagac tactggtaa 780
 tatttttaaa aagtatacag atcaaaagca aaagaagcca tgtgtaacia agagaaatgt 840
 gcaaatattt ttttaaggcag tattaagtgc aagaggagta acatgaaata aacattcttt 900
 cacatggcta ctgggaatat aaatttcgct ccagaaaggc cgtagcagtt tgacgatagg 960
 tggcaaaacc ttaagattgt gtactggggc ccagaatttt tatttctagg aatgtatcct 1020
 gaggaaatta tccgagatcc ccacaaactg caatgttttag gaattgtcct tatagcattg 1080
 catacacaag aaaaacagag aaaagcctga tccctgtcag tggaaaagggt gttcaatgaa 1140
 ttacggtgtg tctgcatgag gcttttatga cattaaaaat tgttgaacia cggccaggca 1200
 cagtggctca tgctgtaat cctaacactt tgggaggcca aggtgggaag attgcctgag 1260
 ctgaggagtt tgagaccagc ctgggcaaca cggtgaaacc ccgtctctac taaaatacaa 1320
 aaaattagcc gggcgctcgca gcatgcgcct gtagtcccag ctgctcagga ggctgaggca 1380
 ggagaattga ttgaacccgg gaggcagagg ttgactgag ctgagattaa gccaccgcac 1440
 tccagcctgg gcgacagagc aagattccgt tcccagaaa aaaaaattgt tcaacaataa 1500
 gggcaaaagg agagaatcat aacatctgat taaacagaaa aagcaagatt tttaaaacta 1560
 actatataag gatggtccca gctgtgtcaa aaggaagctt gtttgtaata cgtgtgcata 1620
 aaaattaaat agaggtgaac acaattattt taaggcagtt aaattatctc tgtattgtga 1680
 actaagactt tctagaattt tacttattca ttctgtactt aaattttttc taatgaacac 1740
 atatactttt gtaatcagaa aatattaaat gcatgtattt ttcaaaatca aaaaaaaaaa 1800
 aaaa 1804

<210> 25
 <211> 2663
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 584050CB1

<400> 25
 ggagaaaggga tggccggcct ggcggcgcggt ttggtcctgc tagctggggc agcggcgctg 60
 gcgagcggct cccaggggcga ccgtgagccg gtgtaccgag actgcgtact gcagtgcgaa 120
 gagcagaact gctctggggg cgctctgaat cacttccgct cccgccagcc aatctacatg 180
 agtctagcag gctggacctg tcgggacgac tgtaagtatg agtgtatgtg ggtcaccgtt 240
 gggctctacc tccaggaagg tcacaaagtg cctcagttcc atggcaagtg gcccttctcc 300
 cggttcctgt tctttcaaga gccggcatcg gccgtggcct cgtttctcaa tggcctggcc 360
 agcctggtga tgctctgccg ctaccgcacc ttcgtgccag cctcctcccc catgtaccac 420
 acctgtgtgg ccttcgcctg ggtgtccctc aatgcatggt tctgggtccac agtctccac 480
 accagggaca ctgacctcac agagaaaatg gactacttct gtgcctccac tgtcatccta 540
 cactcaatct acctgtgctg cgtcaggacc gtgggggctgc agcaccagc tgtggtcagt 600


```

gccttcgagg ctctcctgct gctcatgctg accgtgcacg tctcctacct gagcctcacc 660
cgcttcgact atggctacaa cctggtggcc aacgtggcta ttggcctggt caacgtgggtg 720
tggtggctgg cctggtgcct gtggaaccag cggcggtgc ctcacgtgcg caagtgcgtg 780
gtggtggtct tgctgctgca ggggctgtcc ctgctcgagc tgcttgactt cccaccgctc 840
ttctgggtcc tggatgcca tgccatctgg cacatcagca ccatccctgt ccacgtccctc 900
tttttcagct ttctggaaga tgacagcctg tacctgctga aggaatcaga ggacaagtgc 960
aagctggact gaagaccttg gagcgagtct gccccagtg ggatcctgcc cccgccctgc 1020
tggtccctcc tctccctcca acccttgaga tgattttctc ttttcaactt cttgaacttg 1080
gacatgaagg atgtggggcc agaatacatgt ggccagccca cccctgttg gccctcacca 1140
gccttgaggt ctgttctagg gaaggcctcc cagcatctgg gactcgagag tgggcagccc 1200
ctctacctcc tggagctgaa ctggggtgga actgagtggt ttcttagctc taccgggagg 1260
acagctgcct gtttccctccc caccagcctc ctccccacat cccagctgc ctggctgggt 1320
cctgaagccc tctgtctacc tgggagacca gggaccacag gccttaggga tacagggggg 1380
ccccttctgt taccacccc caccctcctc caggacacca ctagggtgtg ctggatgctt 1440
gttctttggc cagccaaggt tcacggcgat tctcccatg ggatcttgag ggaccaagct 1500
gctgggattg ggaaggagtt tcacctgac cgttgcccta gccaggttcc caggaggcct 1560
caccatactc cctttcaggg ccagggtcc agcaagccca gggcaaggat cctgtgctgc 1620
tgtctgggtg agagcctgcc accgtgtgtc gggagtggtg gccaggtga gtgcataggt 1680
gacagggcgg tgagcatggg cctgggtgtg tgtgagctca ggcctaggtg cgcagtgtgg 1740
agacgggtgt gtccggggaa gaggtgtggc ttcaaagtgt gtgtgtgcag ggggtgggtg 1800
tgtagcgtg ggtagggga acgtgtgtgc gcgtgctggt gggcatgtga gatgagtgc 1860
tgccggtgaa tgtgtccaca gttgagaggt tggagcagga tgagggaatc ctgtcaccat 1920
caataatcac ttgtggagcg ccagctctgc ccaagacgcc acctgggcgg acagccagga 1980
gctctccatg gccaggtgc ctgtgtgat gttccctgtc tgggtccctt ttgcccgcct 2040
cctgcaaac ccacagggtc cccacacaac agtgccctcc agaagcagcc cctcgaggc 2100
agaggaagga aaatggggat ggctggggct ctctccatcc tcttttctc cttgccttcg 2160
catggctggc cttccctccc aaaacctcca tccctctgt gccagcccct ttgccaatgc 2220
ctgatttttg ggaggaggaa ggggcgattt gagggagaag gggagaaagc ttatggctgg 2280
gtctggttcc ttcccttccc agagggtctt aggtttccag ggtggcccca gggcaggcag 2340
gggccacct atgcctgcgc cctggtaaaag gtgacctctg ccatttacca gcagccctgg 2400
catgttctct cccacagga atagaatgga gggagctcca gaaactttcc atcccaaagg 2460
cagttctcgt ggttgaagca gactggattt ttgctctgcc cctgacctct tgctccctct 2520
tgagggaggg gagctatgct aggactccaa cctcagggac tcgggtggcc tgcgctagct 2580
tcttttgata ctgaaaactt ttaagtggtg aggggtggca gggatgtgct taataaatca 2640
attccaagcc tcaaaaaaaa aaa 2663

```

<210> 26
 <211> 769
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 863808CB1

```

<400> 26
gcgacgccga cgcaaggctg ctgctatggg gccggggcgg cgtgtggcgc ggctgctcgc 60
cccactaatg tggcgccagg cggtttcctc ggtggcgggg tccgcggttg gagccgagcc 120
cgggcttcgg ctgctggccg tgcagcggct tcccgtagga gcagcgttct gccgggcttg 180
ccagacccca aactttgtcc gcggcctgca cagcagacct gggctggagg agcggggcga 240
ggggacgggt aacgagggac gccagaatc ggacgggca gatcactatg gtcccaagtt 300
tgacatcgat atgatggttt cacttctgag gcaagaaaat gcaagagaca tttgtgtgat 360
ccaggttcct ccagaaatga gatatacaga ttactttgtg attgttagtg gaacttctac 420
ccgacactta catgccatgg ctttctacgt tgtgaaaatg taaaaacacc tgaatgttaa 480
acgtgacctt catgttaaga tagaaggga ggacactgat gactggctgt gcgtggattt 540
tggcagcatg gtgattcatt tgatgcttcc agaaaccaga gaaatctatg aattagagaa 600
attatggacc ctacgttctt atgatgacca gttagctcag atagcacctg agacagtacc 660
tgaagacttc attcttgga tagaagatga tacttcatct gtgactccag tggagttaaa 720
atgtgaataa aatattttat gcactgcgtt agtcaaaaaa aaaaaaaaaa 769

```

<210> 27
 <211> 1257
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 978433CB1

<400> 27
 gagggcgcg ggtgaaaggc gcattgatgc agcctgcggc ggccctcggag cgcggcggag 60
 cagacgtga ccacgttcct ctctcggtc tcctccgct ccagctccgc gctgcccggc 120
 agccgggagc catgacgccc cagggcccg cgcctcccc gcagcggctc cgcggcctcc 180
 tgctgctcct gctgctgcag ctgcccgcgc cgctcagcgc ctctgagatc cccaagggga 240
 agcaaaaggc gcagctccgg cagaggagg tggtagacct gtataatgga atgtgcttac 300
 aagggccagc aggagtgcct ggtcgagacg ggagccctgg ggccaatggc attccgggta 360
 cacctgggat cccaggctcg gatggattca aaggagaaaa gggggaatgt ctgagggaaa 420
 gctttgagga gtcctggaca cccaactaca agcagtgttc atggagttca ttgaattatg 480
 gcatagatct tgggaaaatt gcggagtgtta catttacaaa gatgcgttca aatagtgttc 540
 taagagtttt gttcagtggc tcacttcggc taaaatgcag aaatgcagtc tgtcagcggt 600
 ggtatttcac attcaatgga gctgaatgtt caggacctct tcccattgaa gctataatgt 660
 atttggaacca aggaagccct gaaatgaatt caacaattaa tattcatcgc acttctctctg 720
 tggaggagct ttgtgaagga attggtgctg gattagtggg tgttgctatc tgggttggca 780
 cttgttcaga ttacccaaaa ggagatgctt ctactggatg gaattcagtt tctcgcacatc 840
 ttattgaaga actacccaaa taaatgcttt aattttcatt tgctacctct ttttttatta 900
 tgccttggaa tgggttactt aaatgacatt ttaaataagt ttatgtatac atctgaatga 960
 aaagcaaagc taaatatgtt tacagaccaa agtgtgattt cacactgttt ttaaactctag 1020
 cattattcat tttgcttcaa tcaaaagtgg tttcaatatt ttttttagtt ggtagaata 1080
 ttttcttcat agtcacattc tctcaacctt taatttggaa tattgttgtg gtcttttgtt 1140
 ttttctctta gtatgcatt tttaaaaaaa tataaaagct accaatcttt gtacaaattg 1200
 taaatgttaa gaattttttt tatatctgtt aaataaaaat tatttccaac aacctta 1257

<210> 28
 <211> 2560
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1655369CB1

<400> 28
 ttccagtga gagcaagtgc tgcccagacc aggacctgt gccaggctag cagccctcca 60
 gctccctcca gagaggaaac ctctgtctgg ctgagggtgg gactagctgg gatgtctcac 120
 tccagtgtct caggttcacc caggaagctc ctccgtggag tggccagcct gattctagcc 180
 ctgtcctctc tggcagcaca tgccacacct gcctgggctt tctgtctcct gatgcttgat 240
 gagccctgc ctctcaatg tttctcaaag acagaccccc ctgaggccag cttgaatgtg 300
 aagactgctg aagtcagctg gcttcaactg agctgcagaa aaggtggctg ggatggccca 360
 ggtgcaccca gagggccag ccttttggct gcctttgggt tgtgacttgg gttgtctctg 420
 aggcctgcc agagctgggc ctgcccgtgg tgggcccgtc gacctcgggc agtcagtgtc 480
 ccgcagcctc agcactgcat cccagaccca gtgtcctcag agggagagc cagcctccct 540
 gctcatgga accaggagtc ccaaaaagtc aggagcctgg aggctctgaa aggagcaggg 600
 attccatagt gcgtgaagct gaaataggcg ccctcctggg gagccccag caaaactgtt 660
 tttcataccc actcccagaa ctgcccgcct ccagctccag cgccagcgcc agctggttgc 720
 caggcgtcat tggagaggcc tggtgcccc aggggcagca gggagtgggt gacctgtatg 780
 ggctggcagg aggcattgg ccattgctgac aagtgtcacc tgccttccca gcctggagcc 840
 accctcagg tggcctgctt gcacctccta tccggaggta gcctgcccc cctgtaggca 900
 gagggggctc ttgcttgagg cctgcacagg aagcaagtat agccccggtg cccagagtg 960
 ggttccactt agccctggcg agatggcctg tcctgagatc tctgtctcca gacccacca 1020
 tctggggagc acagtcctta ggctgcctgg tccaggaagg ggggtgcggt ctgtcaggaa 1080
 acctggactc tcaaggccca ccagcctctc cgtgagtgtt agaaatcaca gatacagtat 1140
 atacttaatt acactaaatt attgctggga ttctcttataa gcactaatta tacctgatta 1200
 taggttaaaa tattttattt gtcaaaatat tttcttggga atgtgtttaa cctttctctg 1260
 gttcattgtt gctgagatgt gaaaactaac cattccctcc tgcctacctt tttggccact 1320
 gggcggcaga gaatggcgct atgtgcagtt gggccctgg caccatgggc ctttggcctg 1380
 cctgctgcag agtagccctg cctgggcagt ctccaggcac tgagcaggcc atctgtggcc 1440
 aggtgagag aatgactggc tcgcttacca gcgtgcagtg gacaaggagc tttggagcct 1500
 caaggggttg ttgctggcct gggctagagg gaaaggtgac catccgtctg tctcctctgc 1560
 tttctattag cgcctccatg tgagtgtatg tgccttgggt cactagcctt cccccaccac 1620
 cccaccatgc cacctggtgg tcttggggcc tgtgtgtgca ctccagcccc tggggaggag 1680
 aggaccagc ccggagagtt ggggcaaggc ctccacatgg cccaagggca acagatgtc 1740

gcagggcagc	tgctgccgat	gctcacgctc	ctgccccct	ccttcccgt	gccacacccc	1800
accctgggccc	cccgcagaca	cgcactctcta	actcagttgg	gcccagcctt	ctggatggct	1860
tggggtaggc	catgggcccc	cctggggcca	ggccagcccc	tggggcagct	ctggaagagc	1920
agtgtggagg	agcacttgct	tgcagcctgg	cttcagcctc	tggcactgct	ggagtggctc	1980
ctgggagctt	ctgcactgtc	ggctttgggg	acgtctcacc	cacttgggtt	acagtaggcc	2040
ttccccaccc	agagagaagt	gtttccaccc	cagagacatt	gtctgtcagc	ccctgaagtg	2100
ctcgctctcc	ccagtgcctg	tcaccagccc	ttcctatctg	tggggtccaa	gtcaggcttc	2160
ccctgcggcc	accagccata	gggagcagcc	atcagccccc	gagtcagaac	tgcttctgtc	2220
tgtccatacc	tccaggtctt	cccggagagg	gggacggata	tttatttctt	aaagtgtgca	2280
cttaattgtg	aggattctca	ggattgttgg	gggctactga	aaagaggaat	gtgttgaatg	2340
tcgctgttgc	tgtccactcg	tcctagaagt	ttagtgtttt	tgctactgtc	atgtgtttct	2400
gtgggcagag	ctggttctgg	aggggtgggtc	agtgaccccg	aggctcagag	catccatcca	2460
ccccactggc	cctccttcca	gataccctct	ctctaattgg	gttcttgcac	gtaaaatact	2520
ccacaataaa	taaataattg	aacaaattaa	aaaaaaaaaa			2560

<210> 29
 <211> 614
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1703244CB1

<400> 29	
gtgcaagagg	aacaaagaaa
tatctcatcc	ttgctggctt
acagatgcct	gctttgtcta
taccctaagg	tgcagatgct
gcctatgtct	tcaccttccc
ggaggcatcg	gccaggcaca
ttcacctacc	gtgtgcctga
gcgctgggcc	cccacctgct
ccaccaccct	ccgacccctt
aggacccagg	actctgttta
gttttagaga	cagg
gggactcctg	cagcgtccgg
cttcactctg	ttccggggcc
tatctaccag	tatgagccat
gatgtacatg	ttttatgtcc
tggttgctcc	tggttccag
atgggggctt	ccatgcacct
ggctgcttct	tcgtgtgcaa
tgccctcagt	ggcccgcat
agccctccac	aagaagcagc
cgtgcccagt	cagccctacc
ctgacctggc	ccttgctcata
tggtgggtgct	tgattgcccc
acctgcggga	ccctgtggcc
tgcccttctg	cgccctggct
actgggcctt	gggtgttgc
gcgcacaccc	gctgtgtat
cttcaccagg	ggcccgcat
ctgtggactc	attgagagag
gggggaagcg	gggggtgggt
	614

<210> 30
 <211> 1936
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1730819CB1

<400> 30	
gactacgggc	tcacagccgt
cgctggggca	acccggctgc
ccggggcagc	ggctgccggg
gggcgaggcc	tgtggcacgg
tgccaacttc	cggaaagcgg
gtcacagcgg	cagctcagcg
tgccctgtac	cgggtccgga
tggtctatgtc	tctcctttg
gctgaccctg	cagctggatg
cgggggctgc	cggggccatg
gcagctgcag	ccgcccacca
ggagatggaa	caggcccaga
atactggatg	tacatcattc
cgggggccag	gggtgggggtg
caggctgggtc	agcgtcccgt
gtgtcctcag	ccatcccaag
ccacctgggc	cagcccttgc
cctttggcac	agcagccggg
gtgggaagaa	gcccagatgg
gctgatggcg	gtagcagcgc
gctgggaggg	gctggggcgg
gctggagcac	tcatttgaga
ctggaaccag	caggatggta
gggccgactc	cgggatgtgg
acccggggcc	ctggatggcc
ctccctgggtg	gagtcgcacc
gtgtcggtgg	gtgtcggtgg
cgtggacctg	gagctgttca
ggcccgactc	gcggccttca
ccctgagacg	cagaagtcct
gttctctcatg	atgtcaggag
gggtgggtgg	ggtagtggcc
caggggcctc	cctttctgct
tttccccccg	tcccacgagg
gaggagcagc	tggactgggg
ccatggcccc	atgcagcccc
	1080

```

aggggcttcc cccctgcccc tggagtagag cccgagatcc tggccactat gccagttctg 1140
acctcgcac cccctacccc gagcccatgc agtctgggaa catgccgcct tctctccagc 1200
ctctgtgcct ttgttccagg tggctcacc ctctgtcccc tggctgggct aggtgggcct 1260
gtccaggctc ctgcagcgcc cccctcactt tgacactgga ctaggatgca gcctcccttc 1320
tgtgtcccc tgagggtacc ctgggtcccc tcatcagggg cagaggcatg aaagagtcgg 1380
ggctggatgg cggggggctt ctgggccccg cgcctagtgc agcccctggg gtcgtggttt 1440
gacatttgct tgcctgggtg aaacaaggaa tccttgccct taaggtgaca ggccctccac 1500
aggcttccag acttgaagga aaaggtttaa gaaagaaaac aaaaccaaca gttagtggag 1560
tcaaagcccc gacactgtaa atagaacccc ctccaccacc ccccgccgcc cagcctccta 1620
cctggactgc ggtgctacga gggcctgcgg gcctttgctg tgtgccacc ccctgtaag 1680
tctattttaa aacatcgacg atacattgaa atgtgtgaac gttttgaaaa gctacagctt 1740
ccagcagcca aaagcaactg ttgttttggc aagacggtcc tgatgtacaa gcttgattga 1800
aattcactgc tcacttgata cgttattcag aaaccaagg aatggctgtc cccatcctca 1860
tgtggctgtg tggagctcag ctgtgttgtg tggcagttta ttaaactgtc cccagatcg 1920
acacgcaaaa aaaaaa

```

<210> 31

<211> 1958

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1757161CB1

<400> 31

```

gccgcgcctt cagctacggc ccgagcgagc ccgcccgcgc cgggcccggc cacagcctgc 60
agcggagccc acgagaggca gcgccatggc ggagcagacc tactcgtggg cctattccct 120
ggtggattcc agtcaagtgt ctacatttct gatttccatt cttcttatag tctatggtag 180
tttcaggctc cttaatatgg actttgaaaa tcaagataag gagaaagaca gtaatagttc 240
ttctgggtct ttcaatggca acagcaccaa taatagcatc caaacaattg actctaccca 300
ggctctgttc cttccaattg gagcatctgt ctctctttta gtaatgttct tcttctttga 360
tcagtttcaa gtagttttta caatatgtac agcagttctt gcaacgatag cttttgcttt 420
tcttctctc ccgatgtgcc agtatttaac aagaccctgc tcacctcaga acaagatttc 480
ctttggttgc tgtggacgtt tcaactgtgc tgagttgctg tcattctctc tgtctgtcat 540
gctcgtctc atctgggttc tcaactggca ttggcttctc atggatgcac tggccatggg 600
cctctgtgtc gccatgatcg cctttgtccg cctgcccagc ctcaaggtct cctgcctgct 660
ttctcaggg cttctcatct atgatgtctt ttgggtattt ttctcagcct acatcttcaa 720
tagcaacgtc atggtgaagg tggccactca gccggctgac aatccccctg acgttctatc 780
ccggaagctc cacctggggc ccaatgttgg gcgtgatgtt cctcgcctgt ctctgcctgg 840
aaaactggct tccccaaagt ccactggcag ccacttctcc atgttgggca tcggagacat 900
cgttatgcct ggtctcctac tatgctttgt ccttcgctat gacaactaca aaaagcaagc 960
cagtggggac tctgtgggg cccctggacc tgccaacatc tccggggcgca tgcagaaggt 1020
ctcctacttt cactgcaccc tcateggata cttttagggc ctgctcactg ctactgtggc 1080
gtctcgcatt caccggggcc cccagcccgc ccttctctat ttggtgccat ttactttatt 1140
gccactcctc acgatggcct atttaaaggg cgacctccgg cggatgtggt ctgagccttt 1200
ccactccaag tccagcagct cccgattcct ggaagtatga tggatcacgt ggaaagtgc 1260
cagatggccg tcatagtcct tttcttcaa ctcatggttt gtttctctt agagctggcc 1320
tggtactcag aaatgtacct gtgtttaagg aactgccgtg tgactggatt tggcatttaa 1380
agggagctcg tttgcaggag agaggtgctg gagccctgtt tggttccttc tcttctgcg 1440
gatgtagagg tggggccctt tccaagaggg acaggcctct cccagcgcg ccttctctcc 1500
acgtttttat ggatctgcac cagactgtta ccttctgggg gagatggaga tttgactgtt 1560
taaaaaactg aacacgcgag gagtctttct agaacttttg aacactaaaa ggatgaaaaa 1620
aattagcaaa ccgaagtttc ttcaatgacc cctcgagaaac tttgggacca gtttctctatg 1680
ggggactcag tttcagagaa ctgagacaga agctcttctg tcgttatatt cttctttcct 1740
ttttttggat ttattaaata ttttctgtgg tgtgaagtga cttattaaat ccacagacat 1800
tgagtgaact cttacaacat ccacataaga atttgttgta atgagttcat gtccacccag 1860
atgttgtgtt ggcagtgaac aagggcacgg tttttatata tacgtacata tatatatata 1920
aacacacaca tagatatata tgaataaaca aaaatgat

```

<210> 32

<211> 1424

<212> DNA

<213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1976095CB1

<400> 32
 gtcaagtagg agacaaggag caaagtccta tcacagcggg aggggacgcc agcgcctgca 60
 gaggtgagc agggaaaaag ccagtgtccc agcgggaagca cagctcagag ctggtctgcc 120
 atggacatcc tgggtcccact cctgcagctg ctggtgctgc ttcttaccct gcccctgcac 180
 ctcatggctc tgctgggctg ctggcagccc ctgtgcaaaa gctacttccc ctacctgatg 240
 gccgtgtgca ctcccaagag caaccgcaag atggagagca agaaacggga gctcttcagc 300
 cagataaagg ggcttacagg agcctccggg aaagtggccc tactggagct gggctgcgga 360
 accggagcca actttcagtt ctaccacccg ggctgcaggg tcacctgcct agacccaaat 420
 cccacttttg agaagttcct gacaaaagagc atggctgaga acaggcacct ccaatatgag 480
 cggtttgtgg tggctcctgg agaggacatg agacagctgg ctgatggctc catggatgtg 540
 gtggtctgca ctctgggtgct gtgctctgtg cagagcccaa ggaaggtcct gcaggaggtc 600
 cggagagtac tgagaccggg aggtgtgctc tttttctggg agcatgtggc agaaccatat 660
 ggaagctggg ccttcatgtg gcagcaagtt ttcgagccca cctggaaaca cattggggat 720
 ggctgctgcc tcaccagaga gacctggaag gatcttgaga acgcccagtt ctccgaaatc 780
 caaatggaac gacagccccc tcccttgaag tggctacctg ttgggccccca catcatggga 840
 aaggctgtca aataatcttt cccaagctcc aaggcactca ttgtctcctt cccagcctc 900
 caattagaac aagccaccca ccagcctatc tatcttcac tgagagggac ctagcagaat 960
 gagagaagac attcatgtac cacctcctag tccctctctc cccaacctct gccagggcaa 1020
 tctctaactt caatcccgcc ttcgacagtg aaaaagctct acttctacgc tgaccaggg 1080
 aggaaacact aggaccctgt tgtatcctca actgcaagtt tctggactag tctcccaacg 1140
 ttgctctccc aatgttgctc ctttccctcg tccccatggg aaagctcctc tcgctttcct 1200
 cctgaggcta caccatgctg tctctaggaa ctggctcaca aagtcatggg gcctgcaccc 1260
 ctgccaagcc cccctgaccc tctctcccca ctaccacctt cttcctgagc tgggggcacc 1320
 agggagaatc agagatgctg gggatgccag agcaagactc aaagaggcag aggttttgtt 1380
 ctcaaatatt ttttaataaa tagacgaaac cacgaaaaaa aaaa 1424

<210> 33
 <211> 2238
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2169991CB1

<400> 33
 cctgtctgtg cacacctcac ggcaagggcc agcctgtttc ctcccgttca cctccaaatc 60
 ttgctgcttt taattcaact cagaggtatg cacttgaggt aggagggcag gggaggtggg 120
 gatggcagga catggatggc ccttgaggca ttggctctgg gtgtcatggg ctgtgagagt 180
 caagaagggc agtggcctgc ctgacttggg ttcgaaaggg tcactctggc cactgcggtg 240
 agaactgaaa ctagccaggc caattctcca ttgttccctg tcttccaggt aggagaatat 300
 tagattaggt ctggagtcag aaggtagcag gggctggggg ttgcaggggg atgttgagaa 360
 gaagtgttcc tttggtcagg gtgggaagcc aacaggattt cctggtgcat tggaggtgaa 420
 agggaaagagg cgtgtgtggt tgtgctttgg gcctgagcag ccagaagccg ttgccatcac 480
 ctatgacaag ggagacagcg cctggggcag agcccagtat ggggtgcatt cagggtagat 540
 caggaggtat gagtgggcag tgggcagacc aggtctgggg ttgtggggga ggcctgggtt 600
 gcaggtttaa gcgtgggacc cacgtcagat tgggtgggtgg tcatgcatgc tgggggtggc 660
 ctcaggtcct cctgcttggc ttcccactcc caggggcttt ctctccag atctccttagc 720
 tgggtgaggg gcaggacaga gccctttcct aggggaagccc ggcaccccct gctgtccagg 780
 gaaggggagt ccttctagcc cctgacagct tctctgcccc tcccctggcc tcccaggcc 840
 tagccagggt tgagttctca cccacctgtg ccgccctgcc ttgttacctg gaagcacagc 900
 cttggggact gagcaggccc tcaactgtcac tttaagaagg gaatcagcca ctttgtgtc 960
 accacctctg ggggaaggtg gagaggagag aagggaagtg ctgtttggct gctgacaaca 1020
 tgaagacttc ctgcgatgag aacagaggca caggtgccgg ccctgcagcc cctgacacct 1080
 ggactggagg gggccatggg gcaccggacc ctggctctgc cctgggtgct gctgaccttg 1140
 tgtgtcactg cggggacccc ggaggtgtgg gttcaagttc ggatggaggc caccgagctc 1200
 tcgtccttca ccatccgttg tgggttccct ggggtctggc ccatctccct ggtgactgtg 1260
 agctgggggg gccccaacg tctgggggg accacgtgg ctgtgttgca ccagaaacct 1320
 ggcattccggc aatgggcccc tgctgcccag gcccgctgg aaaccagag cagcatctct 1380
 ctcatcctgg aaggctctgg ggccagcagc ccctgcgcca acaccacct ctgctgcaag 1440
 tttgcgtcct tccctgaggg ctccctggag gcctgtggga gcctcccgcc cagctcagac 1500
 ccagggtctc ctgccccgcc gactcctgcc cccattctgc gggcagacct ggccgggatc 1560

ttgggggtct	caggagtcct	cctctttggc	tgtgtctacc	tccttcacat	gctgcgccga	1620
cataagcacc	gccctgcccc	taggctccag	ccgtcccgc	ccagccccc	ggcaccgaga	1680
gcacgagcat	gggcaccaag	ccaggcctcc	caggctgtct	ttcacgtccc	ttatgccact	1740
atcaacacca	gctgccgccc	agctactttg	gacacagctc	acccccatgg	ggggccgtcc	1800
tgggtggcgt	cactccccc	ccacgctgca	caccggcccc	agggccctgc	cgcctggggc	1860
tcacacacca	tccttcgacg	tggcagcttt	gtctctgttg	agaatggact	ctacgctcag	1920
gcaggggaga	ggcctcctca	cactggtccc	ggcctcactc	ttttccctga	ccctcggggg	1980
cccagggcca	tgggaaggacc	cttaggagtt	cgatgagaga	gaccatgagg	ccactgggct	2040
ttccccctcc	caggcctcct	gggtgtcacc	cccttacttt	aattcttggg	cctccaataa	2100
gtgtcccata	ggtgtctggc	caggccacc	tgctgcggat	gtggtctctg	tgtgtgcgtg	2160
tgtgggcaca	ggtgtgagtg	tgtgagtgac	agttacccca	tttcagtcac	ttcctgctgc	2220
aactaagtca	gcaacgcc					2238

<210> 34
 <211> 536
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2616827CB1

<400> 34						
gcatgaactt	gggggtcagc	atgctgagga	tcctcttcc	cctggatgta	ggaggagctc	60
aagtgtctgg	aacaggcaag	acccttggg	ctgaaattga	tttcaagtac	gccctcatcg	120
ggactgctgt	gggtgtcgcc	atatctgctg	gcttcctggc	cctgaagatc	tgcatgatca	180
ggaggcactt	atttgacgac	gactcttccg	acctgaaaag	cacgcctggg	ggcctcagtg	240
acaccatccc	gctaaagaag	agagcccaaa	ggcgaaacca	caatttctcc	aaaagagatg	300
cacaggtgat	tgagctgtag	gtgagcagtg	acgtgaagag	gggttctagc	cccgtggaaa	360
acagcccatg	gttaacatct	caggatgttc	tgcattcaaa	cacccaaggc	tggtaatgaa	420
ctttcacatg	gactgaatat	tggaggcaaa	taatagaagg	aatagaatat	acagtgcctc	480
tgtcctgaag	gaaaatatca	tgcctcttct	ggaagaaacg	gactgcacag	aggaag	536

<210> 35
 <211> 2177
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2991370CB1

<400> 35						
cgaggaggctc	gaggccagcc	cgggaccggg	gctgggagca	agcaggcggc	ggcgccggcg	60
gcagaggcgg	cagcgagcgc	ccgcttccca	cgcccctagg	cggcggggcc	gagagcgggg	120
ggatggctcc	gagcgctgac	cccggcatgt	ccaggatgtt	accgttcctg	ctgctgctct	180
ggtttctgcc	catcactgag	gggtcccagc	gggctgaacc	catgttcact	gcagtcacca	240
actcagttct	gctcctgac	tatgacagta	atcccaccca	gctcaactat	ggtgtggcag	300
ttactgatgt	ggaccatgat	ggggactttg	agatcgtcgt	ggcgggttac	aatggaccca	360
acctggttct	gaagtatgac	cgggcccaga	agcggctggt	gaacatcgcg	gtcgatgagc	420
gcagctcacc	ctactacgcg	ctgcgggacc	ggcaggggaa	cgccattggg	gtcacagcct	480
gcgacatcga	cggggacggc	cgggaggaga	tctacttcc	caacaccaat	aatgccttct	540
cggggggtggc	cacgtacacc	gacaagttgt	tcaagttccg	caataaccgg	tgggaagaca	600
tcctgagcga	tgaggtcaac	gtggcccgtg	gtgtggccag	cctctttgcc	ggacgctctg	660
tggcctgtgt	ggacagaaaag	ggctctggac	gctactctat	ctacattgcc	aattacgcct	720
acgggtaagt	gggcccctgat	gccctcattg	aaatggacce	tgaggccagt	gacctctccc	780
ggggcattct	ggcgctcaga	gatgtggctg	ctgaggctgg	ggtcagcaaa	tatacagggg	840
gccgaggcgt	cagcgtgggc	cccatacctca	gcagcagtg	ctcggatata	ttctgcgaca	900
atgagaatgg	gcctaacttc	cttttccaca	accggggcga	tggcaccttt	gtggacgctg	960
cggccagtgc	tgggtgtggac	gacccccacc	agcatgggcg	aggtgtcgcc	ctggctgact	1020
tcaaccgtga	tggcaaatgt	gacatcgtct	atggcaactg	gaatggcccc	caccgcctct	1080
atctgcaaat	gagcacccat	gggaaggtcc	gcttcgggga	catcgctcca	cccaagttct	1140
ccatgccctc	ccctgtccgc	acggtcatca	ccgccgactt	tgacaatgac	caggagctgg	1200
agatcttctt	caacaacatt	gcctaccgca	gctcctcagc	caaccgcctc	ttccgcgtca	1260
tccgtagaga	gcacggagac	cccctcatcg	aggagctcaa	tccccggcag	gccttgggagc	1320

ctgagggccg	gggcacaggg	ggtgtggtga	ccgacttcga	cggagacggg	atgctggacc	1380
tcattctgtc	ccatggagag	tccatggctc	agcogctgtc	cgtcttccgg	ggcaatcagg	1440
gcttcaacaa	caactggctg	cgagtgggtc	cacgcacccg	gtttggggcc	tttgccaggg	1500
gagctaaggt	cgtgctctac	accaagaaga	gtggggccca	cctgaggatc	atcgacgggg	1560
gctcaggcta	cctgtgtgag	atggagcccg	tggcacactt	tggcctgggg	aaggatgaag	1620
ccagcagtg	ggaggtgacg	tggccagatg	gcaagatggt	gagccggaac	gtggccagcg	1680
gggagatgaa	ctcagtgtctg	gagatcctct	acccccggga	tgaggacaca	cttcaggacc	1740
cagccccact	ggagtgtggc	caaggattct	cccagcagga	aaatggccat	tgcatggaca	1800
ccaatgaatg	catccagttc	ccattcgtgt	gccctcgaga	caagcccgtg	tgtgtcaaca	1860
cctatgggaag	ctacaggtgc	cggaccaaca	agaagtgcag	tcggggctac	gagcccacacg	1920
aggatggcac	agcctgcgtg	ggctggtgga	gccctgtgtt	gaagatagtg	acaccacaag	1980
ttgggaagag	ccttggtccc	tgaatcactg	aatcactgcc	ttgaatcacc	gcctggaata	2040
cctgttgatc	aggaacactt	acctggaact	tcactgagca	ggatacaaac	ttctattgta	2100
taaagctatt	aatacattaa	gatttggggg	tgtacactta	cataataaat	tccccatttc	2160
tcttgaaaaa	aaaaaaa					2177

<210> 36

<211> 2043

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3031062CB1

<400> 36

cgccacgacg	cagcggggaa	tctgcagtag	gtctgccggc	gatggagtgg	tggttagct	60
cgccgcttcg	gctctggctg	ctgttggtcc	tctcgccctc	agcgcagggc	cgccagaagg	120
agtcagggtc	aaaatggaaa	gtatttatgt	accaaattaa	caggtctttg	gagaattacg	180
aaccatgttc	aagtcaaaac	tgcagctgct	accatggtgt	catagaagag	gatctaactc	240
ctttccgagg	aggcatctcc	aggaagatga	tggcagaggt	agtcagacgg	aagctaggga	300
cccactatca	gatactaag	aacagactgt	accgggaaaa	tgactgcatg	ttcccccaa	360
ggtgtagtg	tggtgagcac	tttattttgg	aagtgatcgg	gcgtctccct	gacatggaga	420
tggtgatcaa	tgtacgagat	tatcctcagg	ttcctaaatg	gatggagcct	gcatcccgag	480
tcttctcctt	cagtaagaca	tcagagtacc	atgatatacat	gtatcctgct	tggacatttt	540
gggaaggggg	acctgctggt	tggccaattt	atcctacagg	tcttggacgg	tgggacctct	600
tcagagaaga	tctggtaagg	tcagcagcac	agtggccatg	gaaaaagaaa	aactctacag	660
catatttccg	aggatcaagg	acaagtcag	aacgagatcc	tctcattctt	ctgtctcgga	720
aaaacccaaa	acttggtgat	gcagaataca	ccaaaaacca	ggcctggaaa	tctatgaaag	780
ataccttagg	aaagccagct	gctaaggatg	tccatcttgt	ggatcactgc	aaatacaagt	840
atctgtttaa	ttttcgaggc	gtactgcaag	tttccgggtt	aaacacctct	tctgtgtgtg	900
ccattattct	aatgagaaa	agaacatact	agtatggaaa	tttcttaagg	gcagggaagtc	960
atgtatttac	agatgtttat	tgagtacctg	ttatatatca	ggactaagct	gctgggatgt	1020
actagaaaat	acgatatttg	tcactctgct	tccatggaaa	ttttagacta	gcataatacc	1080
tggaaacatag	tagatgaaaa	aaaatacatg	tggtttttagt	ggttgaaatt	taagcatttg	1140
caagttatta	tacttatgac	tcacaaattt	acctttcacc	gaattaagcc	aaaaagactt	1200
tatccagaaa	atagttgagt	agagatgaat	acataggatc	ctgcatgatt	taactttctc	1260
tttgcatttg	ggataataca	ctcaacacat	ttctctacca	catttaaaat	ttttatttat	1320
atcctgcctt	cttcccaaaa	ggattttaaag	cagcttacia	aaatgtatag	cagggcgagg	1380
taaaataaag	aaacagacta	tacaacatgc	atactgtgaa	gtcctcttca	cttccaggta	1440
ccatcatatg	gaacccttat	cctcccacca	caaaatgtat	aattctgctt	atatctgtac	1500
ctgtgatctc	tcttctttca	aaaacgtaga	cctttcttct	tatgaaaaaa	acaatccccc	1560
cacatatatt	ctggatggca	tcttcttca	ctttgtctgg	gactttggct	ctgcccactg	1620
tctcatcat	ccttgatctc	aggaccacag	cctttgtcat	gcagacatat	tcagcttcct	1680
cccatctccc	caacctccag	caaaagaaaa	cccttcattt	tcacattccc	cttcagccga	1740
gattgaagat	gaagatgagg	atgagttcaa	ggatgaagac	caggatgagg	acaaggatga	1800
ggatggagtc	tagagcctcc	cagagcctgg	agaggaggcc	tcggtcagcc	actccgtgga	1860
cgtgggccac	ggtgaccac	catgaagtcc	ccactagcca	ctcgattccc	tgtctgtgca	1920
gagttgctgc	acatcacacc	agccccctgc	aagagcagga	gtcaccacag	gctgaatgcc	1980
cacgaggagc	tctgctgaga	ctctcaaggg	agccagtga	agaaatagaa	ataaagcctg	2040
tgc						2043

<210> 37

<211> 1743

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3101617CB1

<400> 37

```

cagcagggtca cagccccctcg aggcgacagc ggccccgccc caccagagca gtggtacagg 60
catggatggg aagaaatgca gcgtatggat gttcctacct cttgtattta ctttgtttac 120
ttcagctgga ttgtggatag tatacttcat agctgtggaa gatgacaaaa ttttaccatt 180
aaattcagct gaaaggaaac ctggtgtgaa gcatgcacca tatataagca ttgcagggtga 240
tgatcctcct gcaagctgtg tgtttagtca agttatgaac atggcagcct tcctagccct 300
tgtggtagct gttctgcgct tcatacaact gaaaccgaag gttttaaacc cgtggctgaa 360
tattagtgga ttggtggctc tgtgtctggc ttccttcgga atgaccttac ttggttaattt 420
tcagctcaca aatgatgaag aaatccataa cgtcggaaact tccttgacct ttggatttgg 480
cacattgacc tgctggatcc aggtcgcgct gacactcaag gtcaacatca agaataagg 540
acggagaggtt ggaattccac gggttattct gtcggcatct atcactctct gtgtggctct 600
ctacttcatc ctcatggccc aaagcatcca catgtatgca gccagggtcc agtggggcct 660
ggctcatgtgc ttctgtctct attttggcac ctttgccgtg gagttccggc attaccgcta 720
tgagattggt tgctctgagt accaggagaa tttcctaagc ttctcagaaa gcctgtcaga 780
agcttctgaa tatcagactg accagggtgt aaccatcagt ttttccttgc tggtagagggtg 840
gggtgtgacag tgggggaggg gccagtagga cactcaca ggacttgaca tagaacctca 900
tttcacacac acacacacac acacattcat ggccacattt gccaaatgag cttttcaggg 960
cgagttatatt ctttaatgaa aaagcacaag cccttatgtg tcgaaataca cgctgttaca 1020
ctgaaaatat atgcacgaca gagcaagaag cttgtgcatg atcacttctt atccgtcccc 1080
ttcccagcac tccctcctct tcccattctc tccacatgtc tcaagcacc taccgagtag 1140
ggcaggccaa atgttccttg ggagtaatgc caactccga cgttgccctc aggtccaaag 1200
ggcttggaac cagctcgtga ggaagtctg aatctggcac taatattctt gagtggataa 1260
tagtgtatca tagaatagga cggaaattgt attgatgtg gacctgtgt cgctgtgga 1320
aaggcatagt tgaagaact tcccacgaa agccccctc atcgttgctt agtggctcggc 1380
tgtgtggatc ccaggagaga catatgccac agactgtgag agcaaagccc gccgtgtgta 1440
tctggacttg atgcactgtg actgagaatg atttccaaat gtgaatatgt gtagggacgt 1500
ggtctatcag gcctggaaca agatgggggc agtgaaggta tggtttagtg tttgctttca 1560
tagtatgcca tgcacaatgt tttatatttc atagtcttct ttaagtaact accatgagtc 1620
tctctaagcc tcatggacaa agatgtagac caaatgcaag agctgagctt gctttgggtt 1680
caaccatgat caaagaaaaa ctgaggtcac ctgcaggctt acgtgggaag ctaagacaat 1740
atc

```

<210> 38

<211> 1306

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3216178CB1

<400> 38

```

ctgcaaagtt cctgtgagcg ctgtcatttt gtcactctgg tttttcagat tcttccccctg 60
gaggctggag tttccaggat gtcaaaatta cctctgcttg ggtgagctat ttcaagcagc 120
tgggatacct gtgtcactcc tgctgtctgc cagtgactgc ccagggtgtc gctggttcct 180
ccccaggagt agggaggaac cagggtgggt ggctgggatg ggtggatatt taaagaccag 240
gccttggacg ctgcagcact tctatctctg ctgtatgcct gctgccacgt ggctgggtcct 300
cctcctcctg ctgtggctga gccttggggg gaagacaggg agctgctccc aaccccagaa 360
cctttgctgt cttgggacgg atcaccactg caagagggga agttgctact gtgatgaatt 420
ctgccatgtg gcaccagact gccaccacga ccacagtgtc ctctgcaacc ctgcttctca 480
gatgaccaag atggtgctgc agatggtgct gaggtggag aaccaccaa gcccgctag 540
gagccaccta gactggatgc agagcatggt gagctccctg caggttctct gagaaggggt 600
ggatggcagc ctgctccttg cttttgtgcc ctccaggccc caaagtcagg gaaccaaag 660
aagaaagggg cgtagctag ggcagagctc cactgcaatg attgttttag gggtaggagc 720
caggattgcc gtctgtggac actgaaattht gaatctcata tacttttgtg acaaaacatt 780
cttctctttt tgttctctc ctaccatcta aaaatgtaga aaacattctt agcctatgag 840
ttgcacaaaa acaggcagtg gccagatttg gccatagac catagtttgc tgacttctgc 900
cctaaatcat cctccatttc tttccttctg tgtccttgtt actgacaaag ccactttccc 960
taaaatgggg tctttccctg tttggtgcca tgaagccaat atgcaaaacc gaaagtgagc 1020
ctcaagcagt gcaggcttta tttggtggcc atggaattga gaagtgagag cttgggtcac 1080

```


aaatcaactt	ttctgctcgt	gagacccagg	aagtcacaga	tacagggcat	ctttagttaa	1140
ggggctgagc	attaaaagca	aggggaggag	tggccagggt	caatggctca	ctcccataaa	1200
cccagaactt	tgggaggcca	aaatgagagg	attgctgaga	ccaggagtcc	gagaccatcc	1260
tggatcaaat	agtgtataac	ccccatctct	acaaaaataa	aaatga		1306

<210> 39
 <211> 851
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3406803CB1

<400> 39		
gggctggcca	cactgcaggg gctgcaggaa gcaaaggatg aaactgatct tttcactaac 60	
cagcataggg	cacaggccaa agaaaaacttt gggtactctc ttgtgagcca gttgaagtta 120	
gacttggtct	ctttggaata ggtcttcttc tgcagaaata aaaacacttg tctgaaagag 180	
gctgtaacta	tctgtgggct gttgggcaag gacacttcag atactggctt tgagctcact 240	
gggtcctggc	ccctcatggc tctgtgatca tctctgctcc acattgcagg ccatgctccc 300	
tggtgggtgt	cagcccagga gccctccttg ggtcctggcc agactcctcc accctcgtgg 360	
gcctgccgca	accagtctag tgccttttct cccatggggg tccctggaat ctcacacacc 420	
ctgcccctac	agagcctgca gtccagggtg ggagtttaacc ctttccactt tcccagagag 480	
ggagaccttg	agtgaggagg aggtcaggaa acgcggggct ggcagcatgg tgggaggagg 540	
tgaaagtacg	atgaccagag ccttgtgtgt gcggctgttg acaaagctga gggatgatg 600	
gggtgctgtg	gcaggggcgt cctgagaaga ccactcctga tggagtgttg ggggaagtgc 660	
aaggcttgga	ggccaatgag ggggctgatg gcacccatcc agtgaggagg ctgtatttgg 720	
aggccctcct	gagtgaggag aaggctatgt ggatgatgag atgaggtaag tgagatggaa 780	
gtggctgtac	gaacagccat tgaggctgag gcaggaggat tgcctaagtc catagcccag 840	
gtgtcgagac	t	851

<210> 40
 <211> 2204
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3468066CB1

<400> 40	
gagcagtcct	tgtctggccc gccccgcctc ggctcgccgc caggggacgc tagtgggtcc 60
agtctccttg	gcaaccactt gctcctcccc ctccgcccct ttaaccttta ggggtgcgcg 120
gtgcagtata	tctcgcgctc tctccctttt cccccccccc ttttcccacc cggggcgctc 180
aggttgggtc	ggaccggaag cgaagatggc gacttctggc gcggcctcgg cggagctggg 240
gatcgggttg	tgcatattcg gcctcttact actggctatt ttggcattct gctggatata 300
tgttcgtaaa	taccaaagtc ggcgggaaaag tgaagttgtc tccaccataa cagcaatttt 360
ttctctagca	attgcactta tcacatcagc acttctacca gtggatata ttttggtttc 420
ttacatgaaa	aatcaaaatg gtacatttaa ggactgggct aatgctaata tcagcagaca 480
gattgaggac	actgtattat acggttacta tactttatat tctgttatat tgttctgtgt 540
gttcttcttg	atcccttttg tctacttcta ttatgaagaa aaggatgatg atgatactag 600
taaatgtact	caaattaaaa cggcactcaa gtatactttg ggatttgttg tgatttgtgc 660
actgcttctt	ttagttgggt cctttgttcc attgaatgtt cccaataaca aaaattctac 720
agagtgaggaa	aaagtgaagt ccctatttga agaacttggg agtagtcatg gtttagctgc 780
attgtcattt	tctatcagtt ctctgacctt gattggaatg ttggcagcta taacttacac 840
agcctatggc	atgtctgcgt tacctttaaa tctgataaaa ggcactagaa gcgctgtcta 900
tgaacgtttg	gaaaacactg aagacattga agaagtagaa caacacattc aaacgattaa 960
atcaaaaagc	aaagatggtc gacctttgcc agcaagggat aaacgcgcct taaaacaatt 1020
tgaagaaagg	ttacgaacac ttaagaagag agagaggcat ttagaattca ttgaaaacag 1080
ctgggtggaca	aaattttgtg gcgctctgcg tccctgaag atcgtctggg gaatttttt 1140
catcttagtt	gcattgtgtg ttgttaatttc tctctcttgg tcaaathtag ataaagctct 1200
tcattcagct	ggaatagatt ctggtttcat aatttttggg gctaacctga gtaatccact 1260
gaatatgctt	ttgcctttac tacaacagtt tttccctctt gattatattc ttataacaat 1320
tattattatg	tactttattt ttacttcaat ggcagggaatt cgaaatattg gcatatgggt 1380
cttttggatt	agattatata aaatcagaag aggtagaacc aggccccaag cactcctttt 1440

```

tctctgcatg atacttctgc ttattgtcct tcacactagc tacatgattt atagtcttgc 1500
cccccaatat gttatgtatg gaagccaaaa ttacttaata gagactaata taacttctga 1560
taatcataaa ggcaattcaa ccctttctgt gccaaagaga tgtgatgcag aagctcctga 1620
agatcagtgat actgttaccc ggacatacct attccttcac aagttctggg tcttcagtg 1680
tgcttactat tttggtaact gggcctttct tggggatttt ttgattggat taattgtatc 1740
ctgttgtaaa gggaagaaat cggttattga aggatagat gaagattcag acataagtga 1800
tgatgagccc tctgtctatt ctgcttgaca gccttctgtc ttaaagggtt tataatgctg 1860
actgaatatc tgttatgcat ttttaaagta ttaactaac attaggattt gctaactagc 1920
tttcatcaaa aatgggagca tggctataag acaactatat tttattatat gttttctgaa 1980
gtaacattgt atcatagatt aacattttaa attaccataa tcatgctatg taaatataag 2040
actactggct ttgtgagggg atgtttgtgc aaaatttttt cctctaattg ataatagtgt 2100
taaattgatt aaaaatcttc cagaattaat attccctttt gtcacttttt gaaaacataa 2160
taaatacatc gtatctgtgc cttaggttct ccaaaaaaaa aaaa 2204

```

<210> 41
 <211> 570
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3592862CB1

```

<400> 41
gcgcgaggagc tcggggagtc ggcgccatga ccccatcgag gcttccttgg ttgcttagct 60
gggtctcggc cacggcggtg agagcggcaa gatcaccctt tctgtgtcat tctctgagga 120
aaacaagttc ttctcaagga ggaaagtctg aacttgtaa acagtccttt aagaagccga 180
agttaccaga agtcggtttt gatgcaccag aggattccca tttagagaaa gaaccactgg 240
aaaaatttcc agatgatgtt aatccagtga ccaaagaaaa aggtggaccc aggggcccag 300
aacctaccgc atatggagat tgggaacgaa aaggacgctg tattgatttt taagtgcgat 360
attctttaac ttcaatatgt ttttctgaat atgtacatct gaattaactt atttctgatt 420
atttcttttc tttatatcct ttatgtcgtg tagtttgtgt aatgtgttta aatatatata 480
tatatatata tatatatata tatatatatg ggggcttagg aagaaaatat gctgctgttaa 540
attaggaaag ggagaccagc ctgaccaata 570

```

<210> 42
 <211> 802
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3669422CB1

```

<400> 42
caggggtcaag gtgaagctgg tgggtgtctg aggcgggtga gtgtcatggg ggagcctggg 60
tgggggctcac actggctctc tctagtccca tgtcgtcgtc ctcttcacga tgccctctcc 120
cttccccagg gatgtctctg tggagctgcc ttttgttctt atgcacccca agccccacga 180
ccacatcccc ctccccagac ccagtcagg tgagcacact acccacccca agccctcaga 240
gggagggcct gaagcagggc cagtggagga aaactggccc ttccagcacc cacccccaca 300
ccccctcttc cgtcccccc agccctctt ccttccccct cactgggaag cttcttcaac 360
caatcccttc acactctctc ccccatcccc ccaagataca cactggaccc tctcttgcgt 420
aatgtgggca ttaatttttt gactgcagct ctgcttctcc agccccgcgc tgggtggcaa 480
gctgtgttca tacctaaatt ttctggaagg ggacagtga aagaggagtg acaggaggga 540
aagggggaga caaaactcct actctcaacc tcaccaaac acctccatt atcactctct 600
ctgcccccat tccttcaaga ggagaccctt tggggacaag gccgtttctt tgtgaggaa 660
aaaaagggtta gaaggggccc cctctctgaa ggccccact ccctgggatg ctacaatcca 720
atgatggaag atggcattag ctacaccacc ctgcgctttc ccgagatgaa cataccacga 780
actggagatg cagagtcctc ag 802

```

<210> 43
 <211> 693
 <212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3688740CB1

<400> 43

```

gttggttttaa tgggattgtg gaagagaatg actccaatat ttggaagtgc tggtagacca 60
atcagccacc gtccaagaac tgcacacatg cttacctgtc tccgtacccc ttcattgagag 120
gcgagcaciaa ctgcacctcc tatgactctg cagttattta ccgtgggttc tgggcagtc 180
tgatgctcct gggggttagt gctgtagtca tgcgaagctt tttgatcatc tgtgcagccc 240
ccttcgccag cctattttctc tacaaagctg ggggaggctc atatattgct gcagatggaa 300
tttcttctct ctgttactca agcctctcaa agtccttatt gtcccagcct ctgcgtgaaa 360
cgtcttcagc catcaatgac atctcactcc ttcaagccct tatgccactg ctgggatgga 420
ccagtcactg gacctgcatt acagtgggct tatattgacc tttacctcta cacttgatata 480
taactcgttt tcccatttag ttgcaagaca cttgggaagc cagaccaagg cttacatttg 540
tgttttaatg tttttcttgt aaatgcttta tgcctaaatg tttctgtact actcttcttt 600
ccaaatcctt attgtttaaa agtttctctc ctactatacc atgccttata aatattgatt 660
gaatgaatgg atgaaatgca tacctgctta tag 693

```

<210> 44

<211> 1212

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3742589CB1

<400> 44

```

ccctcgaggg aacttgccct tctcaaaccat ggccgccacg gcgcctctgg aagggaaccg 60
ctctggggccc cgcctttgat ctctgttggtg gggctggggg atgagagctg caccgcgcgg 120
gacaagtcgc cggcggcgcc cgacggagca gaacagagag catggagctg gagaggatcg 180
tcagtgcagc cctccttgcc tttgtccaga cacacctccc ggaggccgac ctcatgggct 240
tggatgaggt catcttctcc tatgtgcttg gggctcctgga ggacctgggc ccctcggggc 300
catcagagga gaacttcgat atggaggctt tcactgagat gatggaggcc tatgtgcctg 360
gcttcgcccc catccccagg ggcacaatag gggacatgat gcagaagctc tcagggcagc 420
tgagcgatgc caggaacaaa gagaacctgc aaccgcagag ctctgggtgc caaggtcagg 480
tgcccattct cccagagccc ctgcagcggc ccgaaatgct caaagaagag actaggtctt 540
cggctgctgc tgctgcagac acccaagatg aggcaactgg cgctgaggag gagcttctgc 600
caggggtgga tgtactcctg gaggtgttcc ctacctgttc ggtggagcag gccagtgagg 660
tgctggccaa agctcggggg gacttggaag aagctgtgca gatgctggta gagggaaagg 720
aagagggggc tgcagcctgg gagggcccca accaggacct gccagacgc ctcataggcc 780
cccaaaagga tgagctgaag tccttcattc tgcagaagta catgatggtg gatagcgag 840
aggatcagaa gattcaccgg cccatggctc ccaaggaggc cccaagaag ctgatccgat 900
acatcgacaa ccaggtagtg agcaccaaag gggagcgatt caaagatgtg cggaaccctg 960
aggccgagga gatgaaggcc acatacatca acctcaagcc agccagaaag taccgcttcc 1020
attgaggcac tcgccggact ctgcccagca cttctaggct cagatcccag agggatgcag 1080
gagccctata cccctacaca ggggccccct aactcctgtc ccccttctct actccttgc 1140
tccatagtgt taacctactc tcggagctgc ctccatgggc acagtaaagg tggcccaagg 1200
aaaaaaaaaa aa 1212

```

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 September 2000 (08.09.2000)

PCT

(10) International Publication Number
WO 00/52151 A3

(51) International Patent Classification⁷: C12N 15/00,
C07K 14/47, G01N 33/53

[US/US]; 55 Park Belmont Place, San Jose, CA 95136
(US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs
Drive, Hayward, CA 94545 (US).

(21) International Application Number: PCT/US00/05621

(22) International Filing Date: 3 March 2000 (03.03.2000)

(74) Agents: HAMLET-COX, Diana et al.; Incyte Pharma-
ceuticals, Inc., 3160 Porter Drive, Palo Alto, CA 94304
(US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/123,117 5 March 1999 (05.03.1999) US

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,
SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ,
VN, YU, ZA, ZW.

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/123,117 (CIP)
Filed on 5 March 1999 (05.03.1999)

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): INCYTE
PHARMACEUTICALS, INC. [US/US]; 3160 Porter
Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): TANG, Y., Tom
[CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US).
LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA
95054 (US). BAUGHN, Mariah, R. [US/US]; 14244
Santiago Road, San Leandro, CA 94577 (US). YUE,
Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087
(US). AU-YOUNG, Janice [US/US]; 233 Golden Eagle
Lane, Brisbane, CA 94005 (US). LU, Dyung, Aina, M.

Published:

— With international search report.

(88) Date of publication of the international search report:
26 April 2001

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 00/52151 A3

(54) Title: HUMAN SECRETORY PROTEINS

(57) Abstract: The invention provides human secretory proteins (HSECP) and polynucleotides which identify and encode HSECP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of HSECP.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/05621

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/00 C07K14/47 G01N33/53

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 99 22243 A (ENDRESS GREGORY A ; FLORENCE KIMBERLY A (US); HUMAN GENOME SCIENCES) 6 May 1999 (1999-05-06) abstract page 101 -page 102 Seq.Id.No.215	1-23
A	SHIROZU M ET AL: "CHARACTERIZATION OF NOVEL SECRETED AND MEMBRANE PROTEINS ISOLATED BY THE SIGNAL SEQUENCE TRAP METHOD" GENOMICS,US,ACADEMIC PRESS, SAN DIEGO, vol. 37, no. 3, 1 November 1996 (1996-11-01), pages 273-280, XP002054773 ISSN: 0888-7543 the whole document	1-23

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

1 September 2000

Date of mailing of the international search report

05. 1. 01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Panzica, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/05621

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JACOBS K A ET AL: "A genetic selection for isolating cDNAs encoding secreted proteins" GENE: AN INTERNATIONAL JOURNAL ON GENES AND GENOMES,GB,ELSEVIER SCIENCE PUBLISHERS, BARKING, vol. 198, no. 1-2, 1 October 1997 (1997-10-01), pages 289-296, XP004116069 ISSN: 0378-1119 the whole document	1-23
A	WO 98 30696 A (GENETICS INST) 16 July 1998 (1998-07-16) the whole document	1-23
A	WO 98 32853 A (GENETICS INST) 30 July 1998 (1998-07-30) abstract	1-23
A	WO 98 40486 A (GENETICS INST) 17 September 1998 (1998-09-17) abstract	1-23

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 00/05621

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 16, 19 and 22 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-23 (PARTIALLY)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-23 (partially)

An isolated polypeptide having sequence as set forth in Seq.Id.No.1 of the sequence listing and its nucleic acid sequence as set forth in Seq.Id.No.23. Methods of production and uses thereof.

2. Claims: 1-23 (partially)

Inventions 2-22:

Same as for invention no.1 but respectively to each following pair of aminoacid sequences with their respective nucleic acid sequences:

Invention 2: Seq.Id.Nos.2 and 24.

Invention 3: Seq.Id.Nos.3 and 25.

....

Invention 22: Seq.Id Nos.22 and 44.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/05621

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9922243 A	06-05-1999	AU 1273499 A EP 1042674 A AU 1118499 A WO 9921575 A	17-05-1999 11-10-2000 17-05-1999 06-05-1999
WO 9830696 A	16-07-1998	US 5945302 A US 5972652 A AU 2723397 A EP 0907732 A EP 1007660 A JP 2000508172 T AU 5822398 A AU 2558599 A EP 1047713 A WO 9935168 A	31-08-1999 26-10-1999 29-10-1997 14-04-1999 14-06-2000 04-07-2000 03-08-1998 26-07-1999 02-11-2000 15-07-1999
WO 9832853 A	30-07-1998	AU 5828398 A EP 1007661 A	18-08-1998 14-06-2000
WO 9840486 A	17-09-1998	US 5976837 A AU 6702298 A EP 0973890 A	02-11-1999 29-09-1998 26-01-2000